

ACCEPTED

MAR 20 1985

Under the Federal Insect
Fungicide and Rodenticide
Act amended for the purpose
registered under
EPA Reg. No. *10634-MA-1*

BEST AVAILABLE COPY

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Avoid contamination of feed and food. Remove pets, birds and cover fish aquaria before spraying.

STATEMENT OF PRACTICAL TREATMENT
If Swallowed: Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting of anything by mouth to an unconscious person.
If Inhaled: Remove victim to fresh air. Apply artificial respiration if needed.

If on Skin: Remove contaminated clothing and wash affected areas with soap and water.
If in Eyes: Flush with plenty of water. Call a physician if irritation persists.

ENVIRONMENTAL HAZARDS
This product is toxic to fish. Do not apply directly to lakes, ponds, streams, tidal marshes or estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes.
Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given here-with.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ON LIVESTOCK: To protect cattle and horses from horn flies, house flies, mosquitoes and gnats, apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse flies and deer flies, apply at a rate of 2 ounces per adult cow or horse, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection. To control blood-sucking lice, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

ON PETS: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly. Do not spray into eyes or face. Avoid contact with genitalia. Repeat as necessary.

OUTDOOR APPLICATION
ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest (low marshy places, stagnant pools or ponds and shore lines). Apply at a rate of 2.5 pints per acre (0.1 ounce per 1000 square feet).

Using truck-mounted conventional misting equipment such as Rotomist® models 51C, 91 or 100 HT (traveling at 5 m.p.h. to treat a 300 foot swath) calibrate equipment to deliver 0.9 gallons per minute.

INDOOR APPLICATION
The use of this product in food processing establishments should be confined to lime periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment and thoroughly cleaned with potable water before using.

Vacate treated areas and ventilate before reoccupying.
For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control cockroaches, palmetto bugs, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, meal worms, grain mites and caterpillars. Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under

**AQUEOUS READY TO USE
ANIMAL SPRAY II**

EPA REG. NO. 10634-22 EPA EST. NO. 10634-MA-1

A Concentrated PYRENONE® Micro Emulsion
—Clear—Stable—Leaves No Oil- or Objectionable Residues
Effective, Fast Acting Insecticide
For Use in Homes, Institutions, Restaurants
Food Processing Plants, Dairies, Livestock and Outdoors for Mosquito Control
For Control of Fleas and Brown Dog Ticks on Premises And On Pets
Apply with Conventional Mechanical or Compressed Air Equipment

ACTIVE INGREDIENTS	
Pyrethrins	0.1%
*Piperonyl Butoxide, Technical	1.0%
INERT INGREDIENTS	98.9%
	100.0%
*Equivalent to 0.8% (butylcarbitol) (6-propylpiperonyl) ether and 0.2% related compounds.	

**KEEP OUT OF REACH OF CHILDREN
CAUTION**
See side panel for additional cautions

DO NOT TRANSPORT OR STORE UNDER 32 °F.
PYRENONE—Registered Trademark of Fairfield American Corporation
NET CONTENTS: _____

**MANUFACTURED BY
ALPHA CHEMICAL SERVICES, INC.—STRAIGHTON, MASS.**

pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, mouldings, window and door frames and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

SPI
ven
deli
flies
Any
me
ang
rate
FOX
nan
40,
pos
—
Do
Sto
ten
scr
etc
PE:
Per
cor
api
CO
Tr
cor
or

BATCH _____
Printed in U.S.A.