

READ THE ENTIRE LABEL

CAUTION: Avoid prolonged contact or repeated inhalation or contact with skin. Avoid contamination of food or foodstuffs. In case of contact with skin or eyes flush with plenty of water for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking. Harmful if swallowed. If swallowed, induce vomiting. Call a Doctor.

DIRECTIONS FOR CONTROL OF CLOTHES MOTHS
Sprinkle between layers of clothes being packed away or sprinkle in plastic bags where clothes are being stored.

Use at the rate of 1 ounce for each 5 cubic feet of tightly enclosed space. Will not harm the finest woolsens furs, silks, or other fabric.
Do not mix with naphthalene moth balls or flakes.

Repackaged by

FARM AND RANCH SUPPLY COMPANY

Tulsa, Oklahoma

1 lb. Net Weight



PEACH TREE BORER CONTROL

ACTIVE INGREDIENTS.
PARADICHLOROBENZENE 100%

CAUTION

KEEP OUT OF REACH OF CHILDREN
See Side Panel For Additional Cautions

DIRECTIONS FOR CONTROL OF PEACH TREE BORERS that spend a part of their life cycles in the ground near the trunk.

Distribute the product on the ground in a band 1½ inches from the trunk and 1½ inches wide. Cover the paradichlorobenzene with four inches of soil heaped toward the tree trunk.

If trees under 3 years old are treated, the dosage should be 0.5 oz. per tree. If trees 3 to 6 years old are treated, the dosage should be 0.75 oz. per tree. If trees over 6 years old are treated, the dosage should be 1.0 oz. per tree.

Not recommended for use on trees less than 4 years old unless product is removed at the end of ten days to two weeks.

Fall or Spring applications are the most effective. Fall application is most effective but may be applied in the spring.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

