



# Cowpuncher Fiy Control Number 35

Medicated

## 33 1/3 POUNDS NET WEIGHT

Prevents the breeding of horn flies and face flies and aids in the control of certain gastro-intestinal parasites such as common stomach worms (Haemonchus) lesser stomach worms (Ostertagia), hair worms, bankrupt worms (Trichostrongylus Spp.), hookworms (Bunostomum) and nodular worms (Oesophagostomum) of cattle.

Ethylenediamine Dihydroiodide aids in the prevention of foot rot and soft tissue lumpy jaw in cattle.

PESTICIDALLY ACTIVE DRUG INGREDIENT

Phenothiasine
(6-8 micron average particle size)
1.25%
INERT INGREDIENTS\*
58.75%

Total

100.00%

\*Refers only to ingredients which are neither insecticides nor drugs.

ACTIVE DRUG INGREDIENT
Ethylenediamine Dihydroiodide \_\_\_ 0.032%

## GUARANTEED ANALYSIS

Crude Protein not less than 1	4.0 %
Crude Fat not less than	0.5 %
Crude Fiber not more than	8.0 %
Calcium (Ca) not less than 1	2.00 %
Calcium (Ca) not more than 1	4.00 %
Phosphorus (P) not less than	2.0 %
Iodine (I) not less than	0.02 %
Salt (NaCl) not more than1	5.02 76
Care (1001) not more than 1	5.0 %
Salt (NaCl) not less than1	3.0 %
Vitamin A Dotanom not loss them	
Vitamin A Potency, not less than	ZU,000
USP units per pound.	-

USDA REG. NO. 10640-1

#### INGREDIENTS

Cottonseed meal, calcium carbonate, salcane molasses, grain sorghum gluten feed, dicalcium phosphate, corn gluten feed, soybean meal, iron oxide, steamed bone meal, wheat germ meal, dolomitic limestone, sulphur, vitamin A palmitate (stabilized in gelatin), manganese sulphate, iron sulphate, irradiated dried yeast, copper sulphate, cobalt carbonate and artificial flavor.

WARNING: This feed should be used only in accordance with directions on label.

WARNING: DO NOT TREAT LACTATING DAIRY ANIMALS.

Prevents the breeding of horn flies and face flies but does not kill adult flies. Face flies migrate from farm to farm.

Caution: Consult veterinarian before using in severely debilitated animals. Individual animals are occasionally sensitive to phenothiazine.

#### FEEDING DIRECTIONS

Cowpuncher Fly Control Number 35 Medicated is to be fed free choice. Remove all other mineral and salt. Allow one Cowpuncher Fly Control 35 Block for each five (5) head of cattle. The blocks should be distributed over a large area so as to allow livestock good access to blocks. Cowpuncher Fly Control 35 Medicated eaten at the rate of 6 ounces daily will supply the minimum amount of phenothiazine necessary to prevent the breeding of horn flies and face flies and to aid in the control of certain gastro-intestinal parasites such as common stomach worms, medium stomach worms, hair or bankrupt worms, nodular worms, and hookworms of cattle.

Manufactured By

L AND M MINERAL CO.

Brownwood, Texas

(CAUTION: KEEP OUT OF REACH OF CHILDREN)