

Center Panel

OXY BBC - 12

FOR CONTROL OF

PLANT PARASITIC NEMATODES

Read This Label Carefully

ACTIVE INGREDIENTS

1,2-Dibromo-3-Chloropropane

Other Halogenated C₃ Compounds

INERT INGREDIENTS

TOTAL

BY WT.

81.7%

4.3%

14.0%

100.00%

Oxy BBC 12 contains 12 pounds BBC soil fumigant per gallon.

Patented, U. S. No. 3,049,472. This product licensed for use according to methods of U. S. Patents 2,937,936 and 3,049,472.


EPA Reg. No. 10659-7

WARNING

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed

Harmful liquid and vapor ... Combustible mixture ... May be absorbed through skin ... Causes skin irritation and blisters on prolonged contact ... Do not




Center Panel - Page 2

breathe vapor ... Keep container closed ... Do not take internally ... Wash thoroughly after using ... Do not get in eyes, on skin or on clothing.

Use only in a well-ventilated area. Keep away from heat and open flame, feed and foodstuffs. In case of contact, immediately remove contaminated shoes and clothing and wash skin with soap and water; flush eyes with plenty of water for at least 15 minutes and get medical attention. Do not wear shoes or clothing until completely free of chemical odor. If illness results from inhalation, remove to fresh air and call a doctor. If swallowed, call a doctor. Induce vomiting by giving an emetic such as two tablespoons of salt in a glass of warm water. Do not feed plant remains to dairy or meat animals.

This product is toxic to fish and wildlife, hold any treated water on area to be treated until it is absorbed. Keep out of lakes, ponds and streams. Birds feeding on treated areas may be killed. Apply this product only as specified on this label.


Not for use or storage in or around the home. Rinse equipment and containers and dispose of waste by burying in non-crop lands, away from water supplies. Containers should be destroyed by puncturing and/or smashing and burying with wastes. NEVER REUSE. Do not use containers, pumps or other transfer equipment made of aluminum, magnesium or their alloys, as under certain conditions Oxy BBC 12 may be severely corrosive to such metals.



Manufactured By


OCCIDENTAL CHEMICAL COMPANY
A Division of Occidental Petroleum Corporation
P. O. Box 1185 Houston, Texas 77001

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one, other than an officer of Seller, is authorized to make any other warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage and use of this product is limited to replacement of product or refund of purchase price.



Left and Right Panels

NET 5 GALLONS



GENERAL DIRECTIONS: Apply when soil temperatures at depth of 6 to 8 inches are between 40° and 80° F. The soil should be in good tilth and contain adequate moisture for good seed germination. Extremely wet or dry soils should not be treated. Application on commercial acreages should be made with equipment designed for even distribution at specified depth.

PREPLANT TREATMENT: OVERALL - Apply evenly in rows 12 inches apart at a depth of 6 to 8 inches taking care to seal chisel channels with a drag or press wheel. ROW - Apply in the rows to be planted at a depth of 6 to 8 inches, preferable with two shanks spaced 12 inches apart, placing the fumigant 6 inches to each side of the seed row. Care should be taken to seal chisel channels with a press wheel.

DRENCH TREATMENT: For either pre-plant or post-plant applications in irrigation water, OXY BBC 12 is recommended for use at the rate of 30 ppm of the active ingredient in sufficient water to give penetration to the desired soil depth. Feed the chemical uniformly into the water flow in order to make sure that all the irrigated areas receive equal treatment. One gallon in 48,000 gallons of water is 30 ppm.


NOTE: Any irrigation water treated with OXY BBC 12 MUST BE HELD ON THE TREATED AREA until it is absorbed by the soil. UNDER NO CIRCUMSTANCES IS IT TO BE TURNED INTO THE TAIL WATER OR FED BACK INTO THE IRRIGATION SYSTEM.

MECHANICAL MIXING IS REQUIRED - RECOMMENDED FOR USE ON MINERAL SOILS ONLY.

RECOMMENDED APPLICATION RATES ARE FOR OVERALL TREATMENT. Row application should be made in proportion to overall applications.

DO NOT EXCEED OVERALL PER ACRE RECOMMENDATION IN A SINGLE YEAR.

MIXING DIRECTIONS: This product can be used "as is" or further diluted with mineral spirits. Dilute sufficiently so that the application equipment will deliver the proper amount of soil fumigant per acre. Avoid Spillage When Mixing.



READ THIS LABEL CAREFULLY

RECOMMENDED APPLICATION RATES

VEGETABLES (Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Carrots, Eggplant, Endive, Lettuce, Okra, Parsnips, Radishes and Snap Beans): Pre-plant, at Planting, Postplant - Apply postplant as soon as plants become firmly established, 1½ to 2 gallons per acre.

CUCURBITS (Cucumbers, Melons, Summer Squash): Preplant, at Planting - 1½ to 2 gallons per acre.

TOMATOES: Preplant, Direct Seeded - 1 to 2 gallons per acre. At Planting, Direct Seeded - 1/2 to 3/4 gallon per acre. Preplanting transplants, wait 2 weeks before planting - 1 gallon per acre.

COTTON: Preplant, At Planting, Postplant - 1 to 2 gallons per acre.

SOYBEANS: Preplant, At Planting - ½ to 1 gallon per acre on 42 inch rows.

NOTE: This practice is limited to the following states: Alabama, Arkansas, Delaware, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, South Carolina, Tennessee, Virginia and the southern third of Illinois and Missouri. Do not graze treated areas or feed soybean hay to meat or dairy animals. Do not sell such plant parts to another party or otherwise introduce them into commerce.


PINEAPPLES: Preplant, At Planting, Postplant - Do not apply within 9 months of harvest - 3 to 11 gallons per acre.

BERRIES (Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries): Preplant, At Planting, Postplant - Do not apply within 48 days of harvest; 2 gallons per acre. Strawberries, do not apply within 300 days of harvest; 2 gallons per acre.

CITRUS FRUIT: Preplant, At Planting, Postplant - Do not apply more than once every three years; 3 to 7 gallons per acre.

DECIDUOUS FRUITS (Apricots, Grapes, Nectarines, Peaches): Preplant, At Planting, Postplant - 3 to 7 gallons per acre. Figs: 3 to 5 gallons per acre. Do not apply within one year of harvest.

NUTS: English Walnuts; Preplant, At Planting, Postplant - 5 to 7 gallons per acre. Do not apply within 100 days of harvest.



PEANUTS: Preplant, At Planting - $1\frac{1}{2}$ to 2 gallons per acre. Do not feed,
sell or introduce into commerce hay or hulls taken from treated soil. Forage
crops grown on treated soil should not be used as feed for dairy animals or
animals being finished for slaughter until two years after row treatments or
three years after overall treatments.

