

50 LBS. NET WT.



ZIP

16-5-8

**TURF
FOOD**

with Balan®

9-22-71
10659-10

(Contains 3 pounds Benefin per 250 pounds product)

A Fertilizer-Herbicide Formulation for Use in Turf

GUARANTEED ANALYSIS FOR PLANT FOOD

Total Nitrogen	16%
Available Phosphoric Acid	5%
Potash	8%
Sulfur (Combined with Ammonium Phosphate-Sulfates)	16%
Iron (Complexed in Poly-Phosphoric Acid)	1%
Zinc (Complexed in Poly-Phosphoric Acid)	0.25%

Ingredient Statement for Pest Control

ACTIVE INGREDIENT

N-butyl-N-ethyl-a,a,a-trifluoro- 2,6-dinitro-p-toluidine	1.2%
---	------

INERT INGREDIENTS

Relative to herbicidal content	98.8%
--	-------

®Balan—The trademark for Benefin—Eli Lilly & Company
A division of Eli Lilly & Company

USDA Reg. No. 10659-10

9-22-71
10659-10

CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or absorbed through skin. Avoid prolonged or repeated contact with skin or inhalation of dust. Wash thoroughly after handling. This product is toxic to fish and livestock. Keep out of lakes, ponds and streams. Do not apply where run-off is likely to occur. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes. Birds feeding on treated areas may be harmed. DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.

DIRECTIONS FOR USE:

Do not use this product for any uses other than those specified on the label. ZIPP TURF FOOD is formulated to promote good turf growth and provide Crabgrass and Poa annua (annual bluegrass) control in home lawns, golf courses and parks.

Crabgrass control: For warm season turf grasses, such as bermudagrass, St. Augustinegrass, zoysiagrass, centipedegrass and bahiagrass, apply the entire bag contents to 8712 sq. ft. of area to be treated. BE SURE TO APPLY IN SPRING BEFORE CRABGRASS GERMINATES. For uniform application, set spreader for $\frac{1}{2}$ the recommended rate and apply one-half the total amount over the area to be treated, moving in one direction at a steady rate. Spread the remaining half quantity over the same area, this time moving at right angles to the first application.

Poa annua (annual bluegrass) control: For warm season turf grasses (see above), apply the entire bag contents to 8712 sq. ft. This should be done in late summer or early fall BEFORE annual bluegrass germination. DO NOT APPLY IN THE FALL FOR 4 TO 6 MONTHS FOLLOWING THE SPRING APPLICATION FOR CRABGRASS CONTROL. Application method is the same as for crabgrass control.

Other weeds controlled are barnyard grass (sometimes called watergrass), green and yellow foxtail. Established weeds are NOT CONTROLLED. CHECK SPREADER TO MAKE SURE RECOMMENDED RATE IS BEING APPLIED.

Local conditions affect the use of pesticides. Agricultural experiment stations or extension service agents in the state issue recommendations to fit local conditions and their recommendations should be followed with ZIPP TURF FOOD with Balan.

USE PRECAUTIONS: Do not apply to putting greens. Do not apply to bentgrass. Do not apply to new lawns started the previous fall. Do not apply to dichondra turf or newly sprigged areas of Bermuda grass, St. Augustine grass or centipede grass until these turfs are well established, otherwise root inhibition may result. Balan may thin fine-leaved fescues at rates above those recommended on label. Reseeding should be delayed for at least 6 weeks after applying ZIPP 16-5-8 TURF FOOD with BALAN.

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one, other than an office of Seller is authorized to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond the Seller's control. Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.



Occidental Chemical Company

HOUSTON, TEXAS 77001

