

ACCEPTED
 4-30-73
 1065932

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid inhaling dust or contact with skin, eyes and clothing. Do not use this product where desirable clovers are present or allow the product to "wind drift" on to flowers, ornamental shrubbery or vegetables. Do not apply to newly seeded lawns. During application close spreader openings when turning. Apply when the air is calm. Do not store with other fertilizers, fungicides, insecticides or seeds. Do not apply in areas underlaid by roots of desirable shrubs and trees. Do not reuse container. Destroy by burying in a safe place. Do not burn.

DIRECTIONS FOR USE

(Do not apply this product for any uses other than those specified on the label.)

ZIPP 15-5-10 is a complete turf food containing sulfur, magnesium, iron and zinc in addition to the primary nutrients formulated in a manner that promotes optimum turf nutrition for a longer period of time than that obtained from conventional fertilizers.

For uniform application set spreader for 1/2 the recommended rate and apply one half the total amount over the area to be treated, moving in one direction at a steady rate. Spread the remaining half quantity over the same area, this time moving at right angles to the first application. This product may be used on such turf species as St. Augustine, bahia, bent, Bermuda, centipede and Zoysia. Under conditions of adequate moisture, high temperatures and high humidity, over application may cause some short-lived discoloration of the St. Augustine.

WHEN TO APPLY: For optimum weed control, application should be made when weeds are growing vigorously. Application can be made at any time, however, for maximum safety to the St. Augustine, make application during spring and fall when temperatures are moderate.

AMOUNT TO USE: Apply 50 pounds per 3,000 square feet. Make application when grass blades are dry. CHECK SPREADER FREQUENTLY TO BE SURE THE RECOMMENDED RATE IS BEING APPLIED. For best weed killing results do not mow, rake or water the lawn for at least 24 hours after application. Lawns which are in poor condition (badly infested with weeds) may require a second application of this product. It is suggested this be done one month after the first.

When applied as directed, the following weeds will be controlled:

- | | | | | |
|-------------|---------------|---------------|------------------|--------------|
| Bedstraw | Dandelion | Lespedeza | Poison Ivy | Spurge |
| Black Medic | Dock | Mallow | Poison Oak | Thisle |
| Buckhorn | Ground Ivy | Morning Glory | Purslane | Wild Carrot |
| Burdock | Heal-all | Oxalis | Ragweed | Wild Garlic |
| Chickory | Henbit | Peppergrass | Sheep Sorrel | Wild Lettuce |
| Chickweed | Knotweed | Pigweed | Shepherd's Purse | Wild Onion |
| Clover | Lambsquarters | Plantains | Speedwell | Yarrow |

EPA Reg. No. 10659-32

CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one, other than an officer of Seller, is authorized to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.



a product of
OCCIDENTAL CHEMICAL COMPANY
 a division of
OCCIDENTAL PETROLEUM CORPORATION
HOUSTON, TEXAS



50 lbs. net wt.

Covers 3,000 sq. ft.