

# Green Gro WEED-N-FEED

**CONTAINS 2,4-D**  
**CAUTION: KEEP OUT OF THE REACH OF CHILDREN AND PETS SEE BACK PANEL FOR ADDITIONAL CAUTIONS.**

**10-6-4**  
**WEED-N-FEED**

**NET WT 40 LBS.**

**AMERICAN CHEMICAL CO., SPRINGFIELD, OHIO**

# Green Gro

## 10-6-4

### WEED-N-FEED

ACCEPTED

APR 12 1968

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND REPLICENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
ED UNDER NO. ... SUBJECT  
TO ATTACHED COMMENTS

#### DOUBLE ACTION

**KILLS** — or controls dandelion, buckhorn, clover and many other broadleaf plants.

**FEEDS** — Supplies three primary plant foods for healthier, greener lawns.

**THIS BAG WILL COVER up to 5000 SQ. FT.**

Spreader setting may vary with age, model and condition of the spreader. Make a trial check on a measured area of 1000 square feet with 8 lbs. of product.

#### SPREADER SETTINGS

Scott	7	Jackson	6½	N. K.	4
Vaughn	6½	Cyclone	5½	Lawn Beauty	6

#### DIRECTIONS

For best results apply when the lawn is damp and temperature is warm. Do not apply for at least three days after mowing. Avoid applications when conditions are hot and dry or temperatures are cold. Two to three applications in the spring and early summer are recommended, with at least 60 days between applications.

Not recommended for use on bent grass, dichondra, Centipede grass, St. Augustine grass, on lawns where clover is present or desired, or on new lawns. Apply on a calm day, avoid drift to or contact with vegetables, flowers or shrubs. Thoroughly clean all equipment after each use. Do not store near other fertilizer, seed, insecticides, or fungicides. Spread evenly and avoid over laping.

**Caution:** Keep out of the reach of children.

Avoid contact with skin, eyes and clothes.

Do not take internally.

For content guarantee see front panel. The guarantee of Manufacturer or seller is only that the material contained herein conforms to the information on the label and is reasonably fit for the use here-

# WEED-N-FEED

1000  
LBS  
NET  
WT  
8  
00  
00  
1

## DOUBLE ACTION

KILLS — or controls dandelion, buckhorn, clover and many other broadleaf plants.

FEEDS — Supplies three primary plant foods for healthier, greener lawns.

**THIS BAG WILL COVER up to 5000 SQ. FT.**

Spreader setting may vary with age, model and condition of the spreader. Make a trial check on a measured area of 1000 square feet with 8 lbs. of product.

### SPREADER SETTINGS

Scott	7	Jackson	6½	N. K.	4
Vaughn	6½	Cyclone	5½	Lawn Beauty	6

### DIRECTIONS

For best results apply when the lawn is damp and temperature is warm. Do not apply for at least three days after mowing. Avoid applications when conditions are hot and dry or temperatures are cold. Two to three applications in the spring and early summer are recommended, with at least 60 days between applications.

Not recommended for use on bent grass, dichondra, Centipede grass, St. Augustine grass, on lawns where clover is present or desired, or on new lawns. Apply on a calm day, avoid drift to or contact with vegetables, flowers or shrubs. Thoroughly clean all equipment after each use. Do not store near other fertilizer, seed, insecticides, or fungicides. Spread evenly and avoid over laping.

**Caution:** Keep out of the reach of children.  
Avoid contact with skin, eyes and clothes.  
Do not take internally.

For content guarantee see front panel. The guarantee of Manufacturer or seller is only that the material contained herein conforms to the Chemical described on the label and is reasonably fit for the use herein described when used in accordance with the direction for use. Manufacturer and seller make no other expressed or implied guarantee, including any other express or implied guarantee of fitness or of merchantability.

**MIN. GUARANTEED  
FERTILIZER ANALYSIS**

TOTAL NITROGEN (N).....	10.00%
AVAILABLE PHOSPHORIC ACID (as P <sub>2</sub> O <sub>5</sub> ).....	6.00%
SOLUBLE POTASH (as K <sub>2</sub> O).....	4.00%

**CONTAIN**  
**CAUTION: KEEP**  
**REACH OF CHILDRE**  
**SEE BACK PANEL FOR**  
**CAUTIONS.**

# Green Gro

# PLANT FOOD

**10**  
**LITE**  
**WEED**

**GUARANTEED HERBICIDE ANALYSIS**

ACTIVE INGREDIENTS	
*DIMETHYLAMINE SALT OF 2,4-DICHLOROPHENOXYACETIC ACID	1.00%
INERT INGREDIENTS (PLANT FOOD)	99.00%
<b>TOTAL</b>	<b>100.00%</b>
*EQUIVALENT TO 2,4-DICHLOROPHENOXYACETIC ACID	0.83%
USDA Reg. No.	

**NET 20 LBS.**

**MANUFACTURED BY OCCIDENTAL CHEMICAL CO., SPRINGFIELD, OHIO**

**MIN. GUARANTEED  
FERTILIZER ANALYSIS**

TOTAL NITROGEN (N)..... 10.00%  
AVAILABLE PHOSPHORIC ACID ( $H_2P_2O_4$ )..... 0.00%  
SOLUBLE POTASH ( $K_2O$ )..... 1.00%

# Gree PLANT

**GUARANTEED HERBICIDE ANALYSIS**

ACTIVE INGREDIENTS  
\*DIMETHYLAMINE SALT OF  
2,4-DICHLOROPHENOXYACETIC ACID  
INERT INGREDIENTS PLANT OIL  
EQUIVALENT TO 2,4-DICHLOROPHENOXYACETIC ACID  
100%  
100%

**NET 40**

MANUFACTURED BY OCCIDENTAL CH

**Green Gro**

**PLANT FOOD**

**NET 40 LBS.**

**W**  
**W**  
OCCIDENTAL CHEMICAL CO., SPRINGFIELD, OHIO

**Gro**

**OOD**

**CONTAINS 2,4-D**  
**CAUTION: KEEP OUT OF THE**  
**REACH OF CHILDREN AND PETS**  
**SEE BACK PANEL FOR ADDITIONAL**  
**CAUTIONS.**

**10-6-4**  
**WEED-N-FEED**

SPRINGFIELD OHIO

# Green Gro

# 10-6-4

# WEED-N-FEED

ACCEPTED  
APR 12 1968  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER  
ED UNDER NO. .... SUBJECT  
TO ATTACHED COMMENTS.

## DOUBLE ACTION

**KILLS** — or controls dandelion, buckhorn, clover and many other broadleaf plants.

**FEEDS** — Supplies three primary plant foods for healthier, greener lawns.

**THIS BAG WILL COVER up to 5000 SQ. FT.**

Spreader setting may vary with age, model and condition of the spreader. Make a trial check on a measured area of 1000 square feet with 8 lbs. of product.

## SPREADER SETTINGS

Scott	7	Jackson	6½	N. K.	4
Vaughn	6½	Cyclone	5½	Lawn Beauty	6

## DIRECTIONS

For best results apply when the lawn is damp and temperature is warm. Do not apply for at least three days after mowing. Avoid applications when conditions are hot and dry or temperatures are cold. Two to three applications in the spring and early summer are recommended, with at least 60 days between applications.

Not recommended for use on bent grass, dichondra, Centipede grass, St. Augustine grass, on lawns where clover is present or desired, or on new lawns. Apply on a calm day, avoid drift to or contact with vegetables, flowers or shrubs. Thoroughly clean all equipment after each use. Do not store near other fertilizer, seed, insecticides, or fungicides. Spread evenly and avoid over laping.

**Caution:** Keep out of the reach of children.  
Avoid contact with skin, eyes and clothes.  
Do not take internally.

For content guarantee see front panel. The guarantee of Manufacturer or seller is only that the material contained herein conforms to the Chemical described on the label and is reasonably fit for the use herein described when used in accordance with the direction for use. Manufacturer and seller make no other expressed or implied guarantee, including any other express or implied guarantee of fitness or of merchantability.