


# Green Gro

## 10-6-4



## Lite Weight - Weed - N - Feed

### DOUBLE ACTION

**KILLS** — or controls dandelion, buckhorn, clover and many other broadleaf plants.

**FEEDS** — Supplies three primary plant foods for healthier, greener lawns.

**THIS BAG WILL COVER up to 5000 SQ. FT.**

Spreader setting may vary with age, model and condition of the spreader. Make a trial check on a measured area of 1000 square feet with 4 lbs. of product.

### SPREADER SETTINGS

Scott	6½	Jackson	6	N. K.	4
Vaughn	6	Cyclone	5	Lawn Beauty	5½

### DIRECTIONS

For best results apply when the lawn is damp and temperature is warm. Do not apply for at least three days after mowing. Avoid applications when conditions are hot and dry or temperatures are cold. Two to three applications in the spring and early summer are recommended, with at least 60 days between applications.

Not recommended for use on bent grass, dichondra, Centipede grass, St. Augustine grass, on lawns where clover is present or desired, or on new lawns. Apply on a calm day, avoid drift to or contact with vegetables, flowers or shrubs, Thoroughly clean all equipment after each use. Do not store near other fertilizer, seed, insecticides, or fungicides. Spread evenly and avoid over laping.

**Caution:** Keep out of the reach of children.

Avoid contact with skin, eyes and clothes.

Do not take internally.

For content guarantee see front panel. The guarantee of Manufacturer or seller is only that the material contained herein conforms to the Chemical described on the label and is reasonably fit for the use herein described when used in accordance with the direction for use. Manufacturer and seller make no other expressed or implied guarantee, including any other express or implied guarantee of fitness or of merchantability.

**S**

... 10.00%  
... 6.00%  
... 4.00%

# Green G

# PLANT FOOD

ANALYSIS	
	1.00%
	99.00%
TOTAL	100.00%
ACID	0.83%

## NET 20 LBS.

MANUFACTURED BY OCCIDENTAL CHEMICAL CO

# Green Gro

# 10-6-4

ACCEPTED

APR 12 1968

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. \_\_\_\_\_ SUBJECT  
TO ATTACHED COMMENTS.

## Lite Weight - Weed - N - Feed

### DOUBLE ACTION

**KILLS** - or controls dandelion, buckhorn, clover and many other broadleaf plants.

**FEEDS** - Supplies three primary plant foods for healthier, greener lawns.

**THIS BAG WILL COVER up to 5000 SQ. FT.**

Spreader setting may vary with age, model and condition of the spreader. Make a trial check on a measured area of 100 square feet with 4 lbs. of product.

### SPREADER SETTINGS

Scott	4 1/2	Jackson	6	N. K.	4
Vaughn	5	Cyclone	5	Lawn Beauty	5 1/2

### DIRECTIONS

For best results apply when the lawn is damp and temperature is warm. Do not apply for at least three days after mowing. Avoid applications when conditions are hot and dry or temperatures are cold. Two to three applications in the spring and early summer are recommended, with at least 60 days between applications.

Not recommended for use on bent grass, dichondra, Centipede grass, St. Augustine grass, on lawns where clover is present or desired, or on new lawns. Apply on a calm day, avoid drift to or contact with vegetables, flowers or shrubs. Thoroughly clean all equipment after each use. Do not store near other fertilizer, seed, insecticides, or fungicides. Spread evenly and avoid overlapping.

**Caution:** Keep out of the reach of children.

Avoid contact with skin, eyes and clothes.

Do not take internally.

For content guarantee see front panel. The guarantee of Manufacturer or seller is only that the material contained herein conforms to the

# 10-6-4

APR 12 1968  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND ROUENTICIDE ACT  
FOR ECONOMIC PESTICIDE REGISTERED  
ED UNDER NO. \_\_\_\_\_ SUBJECT  
TO ATTACHED CONDITIONS

## Lite Weight - Weed - N - Feed

### DOUBLE ACTION

- KILLS** or controls dandelion, buckhorn, clover and many other broadleaf plants
- FEEDS** Supplies three primary plant foods for healthier, greener lawns

**THIS BAG WILL COVER up to 5000 SQ. FT.**

Spreader setting may vary with age, model and condition of the spreader. Make a trial check on a measured area of 1000 square feet with 4 lbs. of product.

### SPREADER SETTINGS

Scott . . . . . 6½	Jackson . . . . . 6	N. K. . . . . 4
Vaughn . . . . . 6	Cyclone . . . . . 5	Lawn Beauty . . . . . 5½

### DIRECTIONS

For best results apply when the lawn is damp and temperature is warm. Do not apply for at least three days after mowing. Avoid applications when conditions are hot and dry or temperatures are cold. Two to three applications in the spring and early summer are recommended, with at least 60 days between applications.

Not recommended for use on bent grass, dichandra, Centipede grass, St. Augustine grass, on lawns where clover is present or desired, or on new lawns. Apply on a calm day, avoid drift to or contact with vegetables, flowers or shrubs. Thoroughly clean all equipment after each use. Do not store near other fertilizer, seed, insecticides, or fungicides. Spread evenly and avoid over laping.

**Caution:** Keep out of the reach of children.  
Avoid contact with skin, eyes and clothes.  
Do not take internally.

For content guarantee see front panel. The guarantee of Manufacturer or seller is only that the material contained herein conforms to the Chemical described on the label and is reasonably fit for the use herein described when used in accordance with the direction for use. Manufacturer and seller make no other expressed or implied guarantee, including any other express or implied guarantee of fitness or of merchantability.