

U.S. BORAX  
A MEMBER OF THE RTZ GROUP

20 MULE TEAM  
KILL-POWER

ACCEPTED

August 9, 1972

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 1624-90

UREABOR®

10659-51

NONSELECTIVE WEED AND GRASS KILLER

**UREABOR** is a nonselective weed and grass killer. It is specifically designed for effective control of perennial grasses such as Johnson-grass, quackgrass, Bermuda grass and Dallis grass on industrial sites and noncropped land.

**UREABOR** has a wide range of efficiency in controlling general vegetation. Apply dry by use of a PCB Spreader or use as a water spray.

**UREABOR** is especially effective in eliminating established vegetation. The borate-chlorate gives quick knock-down of weeds and grasses — the bromacil assures long residual control.

**UREABOR** provides control of vegetation for a season or longer.

**UREABOR** is easy to use. Apply dry in granular form, straight from sack to soil with a handy PCB Spreader; or apply as a water spray. **UREABOR** is completely soluble in water.

**UREABOR** is economical to use. Application rates may be from 1/2 to 3 pounds per 100 square feet, depending on vegetation and residual control desired.

**UREABOR** is packaged in 50 pound multiwall paper bags.

*Refer to directions on the label on reverse side for most efficient use of this outstanding product.*

## 20 MULE TEAM<sup>®</sup> KILL-POWER

# UREADOR<sup>®</sup>

## NONSELECTIVE WEED AND GRASS KILLER

On Industrial Sites or  
Similar Noncropped Land

**DRY APPLICATION** — Ready to use granules  
**SPRAY APPLICATION** — Soluble in water

Single Treatment will give Knock-down of vegetation plus Long-lasting prevention of seedlings and regrowth. Especially effective on Perennial Grasses.

### ACTIVE INGREDIENTS:

Sodium Metaborate Tetrahydrate (Na <sub>2</sub> B <sub>4</sub> O <sub>7</sub> ·4H <sub>2</sub> O) . . . . .	66.5%
Boron Trioxide (B <sub>2</sub> O <sub>3</sub> ) equivalent . . . . .	22.6%
Sodium Chlorate (NaClO <sub>3</sub> ) . . . . .	30.0%
Bromacil (5-bromo-3-sec-butyl-6-methyluracil) . . . . .	1.5%
<b>INERT INGREDIENTS</b> . . . . .	2.0%

U.S. Patents 2,700,604 & 3,227,541  
Other Patents Pending      EPA Reg. No. 1624-90

### CAUTION

**KEEP OUT OF REACH OF CHILDREN**  
Avoid contamination of feed and foodstuffs. Do not leave container where children or animals may take internally. Harmful if swallowed. May cause irritation of eyes, nose, throat and skin. Avoid breathing spray mist. Avoid contact with skin and clothing. Do not store in direct contact with ground or concrete floors. Do not reuse container. Destroy when empty.

# USBORAX

A MEMBER OF THE RTZ GROUP  
LOS ANGELES • NEW YORK

ACCEPTED

Aug 9, 1972

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND MITICIDE ACT  
FOR EC REGISTRATION FILED  
UNDER NO. 1624-90

10659-51 DIRECTIONS FOR USE

**WHAT TO USE FOR** — UREADOR is designed for nonselective kill of weeds and grasses with a single treatment. Use for quick knock-down of existing vegetation as well as long lasting effects in the soil for control of regrowth or new seedlings. Use for general mixtures of weeds and grasses. UREADOR is especially potent on perennial grasses such as Johnson-grass, Bermuda grass, Dallis grass and quackgrass.

**WHERE TO USE** — Use UREADOR on noncropped land such as industrial sites, fence lines, storage yards, road sides and railroads.

**WHEN TO USE** — Use UREADOR any time during the growing season. Weeds and grasses are easier to kill with UREADOR when they are young and growing rapidly.

**HOW TO USE** — Uniform application over the entire area to be cleared of vegetation is essential.

**Dry Application** — UREADOR is ready to use. Apply with a shaker can or with a hand operated spreader such as the convenient PCB Spreader designed for this use. Power spreaders can be used for large areas. The dry granules will become effective in the soil after rainfall or artificial watering.

**Spray Application** — Use spraying equipment with mechanical agitation—up to 3 pounds of UREADOR may be dissolved per gallon of water (see application rates below). Add UREADOR gradually to tank during filling with agitator in operation. Dissolve completely before spraying. For quick knock-down of existing vegetation, a thorough wetting of all parts of the plant is recommended.

For small areas or for spot treatment. UREADOR may be dissolved in water and applied with a sprinkling can.

**HOW MUCH TO USE** — Use the following rates of application under most conditions.

Kind of Weeds	Applied	
	Dry	Spray
<b>ANNUAL</b> weeds and grasses such as: sunflower, kochia, Russian thistle, pig-weed, crabgrass, downy brome	1 lb./100 sq. ft.	1 lb./gal./100 sq. ft.
<b>PERENNIAL</b> weeds and grasses such as: Johnsongrass, Bermuda grass, Dallis grass, quackgrass, Canada thistle, field bindweed, bluegrass	3 lbs./100 sq. ft.	3 lbs./gal./100 sq. ft.
<b>MIXED</b> annual & perennial weeds and grasses	3 lbs./100 sq. ft.	3 lbs./gal./100 sq. ft.

As low as 1/2 lb./100 sq. ft. can be used on annual weeds when young and growing rapidly.

As low as 1 lb./100 sq. ft. can be used on seedlings of perennials when young and growing rapidly.

Approximately 2 1/2 cups of UREADOR equals 1 pound.

### IMPORTANT

This material is not selective in action and may be toxic to all types of vegetation and may render the entire treated area totally or partially unproductive for one or more years. Care should be taken to confine the use or application to the particular area intended to be treated and to avoid its contact with lawns, trees, shrubs, crops and other desirable plants which are not intended to be destroyed or injured. This includes precautions in treating areas which may be underlaid by roots of adjacent valuable growths. Careless application of this material, or washing by water run-off, to areas where desirable plants are growing or which will be used for later planting may result in injury to desirable plants. Do not drain or flush equipment on or near such areas. Do not contaminate domestic or irrigation water or waters used by fish and wildlife. Wash clothing and equipment after application. Keep animals off treated areas until rainfall has leached all the herbicide into the soil. Mechanical equipment should be thoroughly cleaned after use.

### NOTICE

Because of the many varying conditions affecting use and application, such as location, seasonal weather, soil composition and type of vegetation, manufacturer warns buyer that these and other conditions may impair or vary the results or effects of the use of this product. In any event, complete elimination of weeds is not guaranteed. It is advisable to consult your State Extension Service or Agricultural Experimental Station weed specialists on specific problems. Neither manufacturer nor seller shall be liable in respect to any injury or damage suffered by reason of the use of this product for a purpose not indicated by the label or when used contrary to the directions or instructions hereon nor with respect to breach of any warranty not expressly specified herein. Buyer accepts this material subject to these terms, and assumes all risk of usage and handling except when used or handled in accordance with this label. Neither manufacturer nor seller makes, and their employees and agents are not permitted to make, any representations or warranties, express or implied by law, not specifically set forth herein.

**UNITED STATES BORAX & CHEMICAL CORPORATION**

3075 WILSHIRE BLVD., LOS ANGELES, CALIFORNIA 90010