



SENTRY

DIRECTIONS

FOR PROFESSIONAL USE ONLY

TF33

May be fatal if swallowed when spraying. Do not get on skin or clothing after handling.

Sentry TF33 should be used in conjunction with good turf management. For a preventive program, apply every 7 to 10 days starting before diseases normally appear and continuing throughout the season. If a preventive program is not followed, inspect turf daily and apply at the first sign of disease. For the treatment of golf greens, always treat aprons and approaches. If soil is dry, water thoroughly before applying TF33. Avoid treatment during heat of day. preferably use in early morning or late evening. Do not water for 24 hours after treatment. After spraying Sentry TF33, thoroughly clean the spraying equipment with water.

Fungicide for prevention and control of Dollar Spot, Copper Spot, Helminthosporium and Curvularia Leaf Spots in fine turf grasses.

ACTIVE INGREDIENT:

Cadmium Chloride* 20.1%

INERT INGREDIENTS:

. 79.9%

100.0%

EXTERNAL — In case of contact with large volumes of water. If in eyes, wash with water and call a physician. Wash hands with water before reuse.

INTERNAL — Call a physician immediately. If swallowed, drink warm water, and repeat with egg beaten with water.

For Turf grasses: ^(PLC)
Dollar Spot (Sclerotinia homeocarpa), Copper Spot, Leaf Spots (Helminthosporium and Curvularia).

To each 1000 square feet of turf to be treated, apply one (1) ounce of TF33 diluted with 5 to 10 gallons of water. Repeat every 7 to 10 days either until disease conditions are past and control is obtained, or for preventive maintenance.

Keep from Freezing

Storage of this product in cold or freezing temperatures may cause it to thicken or separate. Performance will not be affected. However, before using stir thoroughly and allow product to warm to at least 60°F.

*Equivalent to 12.3% as elemental cadmium.

ACCEPTED
7/19/72
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECOTOXIC POISON REGISTERED UNDER NY 100-11111 SUBJECT TO ATTACHMENT

**DANGER
POISON**



KEEP OUT OF REACH OF CHILDREN

(See additional precautions on right panel)

Sold By

SENTRY CHEMICAL COMPANY • P. O. BOX 28895 • ATLANTA, GA. 30328

FISH AND WILDLIFE — This product is toxic to fish and wildlife. Do not apply to lakes, streams, or ponds. Do not use equipment, or disposal of this label.

DOMESTIC ANIMALS — do not use grass clippings from treated areas for animal bedding and when there is doubt about the safety of the material.

CONTAINER DISPOSAL — Do not reuse container. Do not burn, bury, or crush and dispose of in large quantities of water.



SENTRY



FOR PROFESSIONAL USE ONLY

TF33

Fungicide for prevention and control of Dollar Spot, Copper Spot, Helminthosporium and Curvularia Leaf Spots in fine turf grasses.

ACTIVE INGREDIENT:

Cadmium Chloride* 20.1%

INERT INGREDIENTS: 79.9%

100.0%

*Equivalent to 12.3% as elemental cadmium.

... good turf management.
... 10 days starting before
... oughout the season. If a
... rf daily and apply at the
... eens, always treat aprons
... ly before applying TF33.
... y use in early morning or
... treatment. After spraying
... uipment with water.

... Spot, Leaf Spots (Helmin-

... l, apply one (1) ounce of
... epeat every 7 to 10 days
... ontrol is obtained, or for

... peratures may cause it to
... ffected. However, before
... rm to at least 60°F.

ACCEPTED
7/19/72
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND FUNGICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 10663-7 SUBJECT
TO ATTACHMENT

**DANGER
POISON**



KEEP OUT OF REACH OF CHILDREN

(See additional precautions on right panel)

Sold By

SENTRY CHEMICAL COMPANY • P. O. BOX 28895 • ATLANTA, GA. 30328

DANGER

May be fatal if swallowed. Do not breathe spray mist—wear respirator when spraying. Do not get in eyes, or on skin, or clothing. Wash thoroughly after handling.

ANTIDOTE

EXTERNAL — In case of contact, wash immediately with soap and large volumes of water. If in eyes, flush with water for at least 15 minutes and call a physician. Wash contaminated clothing with soap and hot water before reuse.

INTERNAL — Call a physician at once! If swallowed, give raw white of egg beaten with water; then give a tablespoonful of salt in a glass of warm water, and repeat until vomit fluid is clear. Repeat raw white of egg beaten with water.

FISH AND WILDLIFE — This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DOMESTIC ANIMALS — do not let animals feed or graze on treated areas. Do not use grass clippings for feed. Keep animals away during application and when there is danger from drift or residues.

CONTAINER DISPOSAL — Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place. Flush spillage with large quantities of water.

EPA Reg. No. 10663-7



SENTRY

DIRECTIONS

FOR PROFESSIONAL USE ONLY

TF33

Sentry TF33 should be used in conjunction with good turf management. For a preventive program, apply every 7 to 10 days starting before diseases normally appear and continuing throughout the season. If a preventive program is not followed, inspect turf daily and apply at the first sign of disease. For the treatment of golf greens, always treat aprons and approaches. If soil is dry, water thoroughly before applying TF33. Avoid treatment during heat of day. preferably use in early morning or late evening. Do not water for 24 hours after treatment. After spraying Sentry TF33, thoroughly clean the spraying equipment with water.

For Turf grasses:

Dollar Spot (Sclerotinia homeocarpa), Copper Spot, Leaf Spots (Helminthosporium and Curvularia).

To each 1000 square feet of turf to be treated, apply one (1) ounce of TF33 diluted with 5 to 10 gallons of water. Repeat every 7 to 10 days either until disease conditions are past and control is obtained, or for preventive maintenance.

Keep from Freezing

Storage of this product in cold or freezing temperatures may cause it to thicken or separate. Performance will not be affected. However, before using stir thoroughly and allow product to warm to at least 60°F.

Fungicide for prevention and control of Dollar Spot, Copper Spot, Helminthosporium and Curvularia Leaf Spots in fine turf grasses.

ACTIVE INGREDIENT:

Cadmium Chloride* 20.1%

INERT INGREDIENTS:

. 79.9%

100.0%

*Equivalent to 12.3% as elemental cadmium.

10663-7
 ACCEPTED
 7/19/92
 UNREGISTERED FUNGICIDE
 FOR REGISTERED UNDER NY 10663-7

DANGER

POISON



KEEP OUT OF REACH OF CHILDREN

(See additional precautions on right panel)

Sold By

SENTRY CHEMICAL COMPANY • P. O. BOX 28895 • ATLANTA, GA. 30325

M
W
O
E
V
A
W
I
E
W
E
F
O
E
O
D
D
T
I
C
F
L

SENTRY



FOR PROFESSIONAL USE ONLY

TF33

Fungicide for prevention and control of Dollar Spot, Copper Spot, Helminthosporium and Curvularia Leaf Spots in fine turf grasses.

ACTIVE INGREDIENT:

Cadmium Chloride* 20.1%

INERT INGREDIENTS:

. 79.9%

100.0%

*Equivalent to 12.3% as elemental cadmium.

DANGER

POISON



KEEP OUT OF REACH OF CHILDREN

(See additional precautions on right panel)

Sold By

DANGER

May be fatal if swallowed. Do not breathe spray mist—wear respirator when spraying. Do not get in eyes, or on skin, or clothing. Wash thoroughly after handling.

ANTIDOTE

EXTERNAL — In case of contact, wash immediately with soap and large volumes of water. If in eyes, flush with water for at least 15 minutes and call a physician. Wash contaminated clothing with soap and hot water before reuse.

INTERNAL — Call a physician at once! If swallowed, give raw white of egg beaten with water; then give a tablespoonful of salt in a glass of warm water, and repeat until vomit fluid is clear. Repeat raw white of egg beaten with water.

FISH AND WILDLIFE — This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DOMESTIC ANIMALS — do not let animals feed or graze on treated areas. Do not use grass clippings for feed. Keep animals away during application and when there is danger from drift or residues.

CONTAINER DISPOSAL — Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place. Flush spillage with large quantities of water.

EPA Reg. No. 10663-7

SENTRY CHEMICAL COMPANY • P. O. BOX 28895 • ATLANTA, GA. 30328

10663-7
ACCEPTED
7/19/72
UNL...
FUNGIC...
FOR EC...
ED UNDER NO. 10663-7