

PAL 11 TP 11/21/57



999
ODORLESS • CONCENTRATED
insecticide
spray

ACTIVE INGREDIENTS

ACTIVE INGREDIENTS	PERCENTAGE
Pyrethrin	0.25
Piperonyl Butoxide, Technical	0.52
N-octyl bicycloheptene Dicarbosimide	0.83
Consists of (Butylcarbitol) (6-Propylpiperonyl) ether and other related compound	98.40
Inert ingredients	100.00

CAUTION:

KEEP OUT OF REACH OF CHILDREN
 SEE BACK PANEL FOR ADDITIONAL
 CAUTIONS

EPA REG. NO. 10693-7
 EPA EST. NO. 10693-CA-1
 Manufactured by:

FLO-KEM INC. Phone 213 632-7124 19402 Susana Rd., Compton, Calif. 90221

DIRECTIONS FOR USE

Flies, Mosquitoes, Gnats: Close room and discharge mist vigorously in all parts of room, using 1 to 1 1/2 ounces for each 1000 cubic feet of space. Leave room closed for at least ten minutes. Sweep up and destroy fallen insects.

Fleas (Infesting Premises): Remove and destroy animals' bedding and spray sleeping quarters freely, forcing the spray into all cracks in the floor and about the room. Repeat when re-infested.

Roaches, Waterbugs: Apply in all cracks and crevices in woodwork, underneath sinks, dark places behind pipes, and in all places where roaches and waterbugs infest, contacting as many insects as possible. Repeat operation as often as necessary.

Clothes Moths, Carpet Beetles: Spray all articles to be protected, giving particular attention to folds and seams. Interior surfaces of storage containers such as closets, chests and trunks should be sprayed regularly. The treatment should be repeated every 30 days unless the articles are stored in moth-tight containers.

Grain Weevils, Cadellids, Flour Beetles, Grain Mites (Accessible Stages): Wet surfaces thoroughly and treat all infested surface areas, insect runways, and hiding places at the rate of 1 gallon of this material to 1000 square feet of area, or to the point of runoff. Repeat when necessary. **DO NOT SPRAY STORED GRAIN.**

FOOD PROCESSING PLANTS: Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product as a space spray in these areas, apply only when plant is not in operation. Product should be applied in edible product areas of federally inspected meat, poultry, and egg packing plants only as a space spray.



**DO NOT REMAIN IN TREATED AREA
 VENTILATE AREA WHEN TREATMENT IS COMPLETED**

This spray is especially designed for use in restaurants, food processing plants, food storage warehouses, flour mills, meat packing plants, animal kennels, and the like. (Meat Inspection Division Memorandum No. 52—Insect Sprays.) This spray is designed for use in either fogging equipment or hand sprayers.

CAUTION:

KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Do not spray on floor stuff. Do not pour, store or spill near heat or open flame. Keep containers closed.

MSK 204 Insecticide Synergist



Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

BEST AVAILABLE COPY