



**DISINFECTANT • DEODORIZER • SANITIZER**

**ACTIVE INGREDIENTS:**

n-Alkyl (60% C<sub>14</sub>, 30% C<sub>16</sub>, 5% C<sub>12</sub>, 5% C<sub>18</sub>)  
 dimethyl benzyl ammonium chlorides ..... 5%  
 n-Alkyl (68% C<sub>12</sub>, 32% C<sub>14</sub>) dimethyl  
 ethylbenzyl ammonium chlorides ..... 5%

**INERT INGREDIENTS:** ..... 90%  
 100%

Trophy No-Bac is a concentrate and should be diluted before using.

**DANGER: KEEP OUT OF REACH OF CHILDREN.**

(See other precautionary and first aid statements on left side panel.)

**PURDY PRODUCTS INC.**  
**MILWAUKEE, WISCONSIN 53208**

**WARRANTY**

are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, acts of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must or damage, resulting from its misuse as such, or in combination with other materials. Statements concerning the use of these pending the infringement of any patent, and no liability for infringement arising out of any such use is assumed.

**USE DIRECTIONS:**

**Household** — For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Locker Rooms, Garbage Pails, Sink Tops, Corridors, Classrooms, Offices and shower Stalls. Apply solution with mop or cloth. Sanitize with 1 ounce Trophy No-Bac to 4 gallons of water. Disinfect with 3 ounces Trophy No-Bac to 5 gallons of water. Kills Staphylococcus aureus and Salmonella choleraesuis. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

**Hospitals** — For floors, walls and other hard surfaces. In Nursing Homes, Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms, use Trophy No-Bac at 3½ ounces per 5 gallons of water. Apply with cloth, mop, or sponge. At this level Trophy No-Bac will be effective against Pseudomonas aeruginosa PRD-10. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

**Food Processing Equipment** — For sanitization of previously cleaned food processing equipment and food utensils dilute 1 ounce Trophy No-Bac per 4 gallons of water to provide 200 ppm of active quaternaries. At this level, **No Potable Water Rinse** is required.

**Dairies** — To sanitize dairy equipment such as tanks, lines, pails, and milk cans, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce Trophy No-Bac to 4 gallons of water (200 ppm). At this level, **No Potable Water Rinse** is required. Follow recommendations of local Health Board.

**Restaurant and Bar Rinse** — Dishes, glassware, silverware, cooking utensils, wash with soap or synthetic detergent; rinse thoroughly and immerse in a sanitizing solution containing 1 ounce Trophy No-Bac to 4 gallons of water. **No Terminal Potable Water Rinse Required.**

This product when used on environmental, inanimate, hard surfaces at 3½ ounces per 5 gallons of water is effective against Influenza A<sub>2</sub>, Herpes Simplex, Adenovirus type 5, and Vaccinia Viruses.

NET CONTENTS: 1 GALLON