

04 FEB 1985

04 FEB 1985

Purdy Products, Inc.  
1526 North 31st Street  
Milwaukee, WI 53208

Attention: James P. Kucharski  
Executive Vice President

Gentlemen:

Subject: Trophy No-Bac  
EPA Registration No. 10710-3  
Your Submission Dated October 23, 1984

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable, provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

a. Revise the statement:

Causes eye damage and skin irritation

to read:

Causes severe eye and skin damage

Ref: The Onyx label you referenced

b. Add the statement:

Statement of Practical Treatment

to appear directly below the word: DANGER  
(center panel only)

CONCURRENCES


SYMBOL ▶								
SURNAME ▶								
DATE ▶								

3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

John H. Lee   
Product Manager (31)  
Disinfectants Branch  
Registration Division (TS-767C)

Enclosure

## Precautionary Statements Hazards to Humans and domestic animals

### **DANGER**

Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Do not breathe spray mist. Avoid contamination of food.

### **STORAGE AND DISPOSAL**

- DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL
- OPEN DUMPING IS PROHIBITED
- DO NOT REUSE EMPTY CONTAINER

### **PESTICIDE DISPOSAL**

PESTICIDE SPRAY MIXTURE OR RINSE WATER THAT CANNOT BE USED ACCORDING TO LABEL INSTRUCTIONS MUST BE DISPOSED OF ACCORDING TO FEDERAL OR APPROVED STATE PROCEDURES UNDER SUBTITLE C OF THE RESOURCE CONSERVATION AND RECOVERY ACT.

### **CONTAINER DISPOSAL**

TRIPLE RINSE (OR EQUIVALENT) THEN OFFER FOR RECYCLING OR RECONDITIONING, OR DISPOSE OF IN A SANITARY LANDFILL, OR BY INCINERATION IF ALLOWED BY STATE AND LOCAL AUTHORITIES

### **GENERAL**

CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES

**Trophy No-Bac** provides sanitizing, disinfecting and deodorizing in Hospitals, Food Processing Plants, Dairies, Restaurants, Bars, and other institutions. When used as recommended, it is effective as a sanitizer against *Escherichia coli* and *Staphylococcus aureus*; and as a disinfectant against *Staphylococcus aureus*, *Salmonella choleraesuis* and *Pseudomonas aeruginosa*. Trophy No-Bac is authorized under the Food Additive Amendment of the Federal Food, Drug and Cosmetic Act for use as a sanitizing solution at 200ppm active level without requiring a final potable water rinse.

**DISINFECTION**- At 3 ounces per 5 gallon dilution Trophy No-Bac exhibits effective disinfectant activity against the organisms: *Salmonella choleraesuis* and *Staphylococcus aureus*

At 3 1/2 ounces per 5 gallon dilution, Trophy No-Bac is an acceptable hospital disinfectant and demonstrates disinfectant activity against *Pseudomonas aeruginosa*.

**VIROCIDAL ACTIVITY**- This product when used on environmental, inanimate hard surfaces at 3 1/2 ounces per 5 gallons of water is effective against Influenza A<sub>1</sub> (representative of the flu virus), Herpes simplex (causative agent of fever blisters and mononucleosis), Adenovirus Type 2 (causative agent of upper respiratory infections), and Vaccinia Virus (representative of the pox virus group).

**EFFICACY TESTS HAVE DEMONSTRATED THAT Trophy No-Bac IS AN EFFECTIVE BACTERICIDE, AND VIRUCIDE\* IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).**

**DEODORIZATION**- Trophy No-Bac deodorizes those areas which generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, pet areas and any other areas which are prone to odors caused by microorganisms.

**MOLD & MILDEWSTAT**- Trophy No-Bac prevents and controls mold and mildew and the odors they cause on hard non-porous inanimate surfaces.

**SANITIZATION**- When used for sanitization of previously cleaned food equipment or food contact items limit active quaternaries to 200ppm. At this level **NO POTABLE WATER RINSE** is required

Trophy No-Bac is an effective sanitizer when diluted in water up to 750 ppm hardness (CaCO<sub>3</sub>)

**BEST AVAILABLE COPY**

ACTIVE INC

INERT INC

In case  
for at  
tamina

If swa  
soluti  
Avoid

NOTE T  
use of  
depres

SEE LE

EPA Est

## DIRECTIONS FOR USE

### GENERAL CLASSIFICATION

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

Trophy No-Bac has been designed specifically for hospitals, food processing plants, dairies, restaurants, bars, and other institutions where disinfection and sanitization is of prime importance

**DISINFECTION**- To disinfect previously cleaned hard inanimate surfaces (such as walls, floors, table tops), apply Trophy No-Bac with mop, cloth, or sponge or mechanical sprayer so as to wet thoroughly. Allow to remain wet for 10 minutes and then let air dry

**DILUTION RATES**- General Disinfection - add 3 ounces per 5 gallons of water. Hospital Disinfection - add 3 1/2 ounces per 5 gallons of water.

For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1 1/2 ounces Trophy No-Bac per gallon of water for 10 minutes.

For disinfection, a pre-cleaning step is required and recommended. Prepare a fresh solution for each use

**DEODORIZER**- To deodorize, apply Trophy No-Bac as indicated under the heading "General Disinfection" above.

**MILDEWSTAT**- To control mold and mildew on pre-cleaned, hard non-porous surfaces (such as floors, walls, table tops), add 3 ounces Trophy No-Bac per gallon of water. Apply solution with a cloth, mop, or sponge making sure to wet all surfaces completely. Let air dry. Prepare a fresh solution for each use. Repeat application at weekly intervals or when mildew growth reappears.

**SANITIZATION**- To sanitize previously cleaned and rinsed non-porous food contact surfaces, prepare 200ppm active quaternary solution by adding 1 ounce Trophy No-Bac to 4 gallons of water

To sanitize immobile items (such as tanks, chopping blocks, counter tops), flood the area with 200 ppm Trophy No-Bac for at least 60 seconds making sure to wet all surfaces completely. Remove and let air dry. Prepare a fresh solution daily or more frequently as soil is apparent.

To sanitize mobile items (such as drinking glasses, eating utensils), immerse in 200ppm Trophy No-Bac for at least 60 seconds making sure to immerse completely. Remove and let air dry. Prepare a fresh solution daily or more frequently as soil is apparent.

### STATE OF WISCONSIN FIVE STEP SANITIZATION PROCEDURE

1. Scrap and prewash utensils and glasses whenever possible.
2. Wash with a good detergent or compatible cleaner (a commercial cleaner or detergent can be recommended as long as it is not of an anionic nature).
3. Rinse with clean water.
4. Sanitize in a solution of 1 ounce to 4 gallons of water (200ppm). Immerse all utensils for at least 2 minutes or contact time specified by governing sanitary code.
5. Place sanitized utensils on racks or drainboard to air dry.

NOTE: A clean potable water rinse following sanitization is not permitted under Section H 96.06 (3) (h) of the Wisconsin Administrative code.

### DISINFECTION OF POULTRY EQUIPMENT ANIMAL QUARTERS AND KENNELS

Brooders, founts, feeding equipment and animal quarters (stalls, etc.) can be disinfected after cleaning by applying a solution of 1 ounce Trophy No-Bac to 5 gallons of water. Sanitized utensils should be immersed in solution.

Prior to disinfection, all poultry animals, and feeds must be removed from area. Empty all troughs, feeding, and watering appliances. Remove all litter and droppings from floors, walls and surfaces occupied or traversed by poultry or animals.

After disinfection, ventilate brooders, coops and closed spaces. Do not house poultry, animals, or equipment until treatment has been absorbed, set or dried.

All treated equipment that comes in contact with feed or drinking water must be rinsed with potable water before reuse.

**BEST AVAILABLE COPY**