

# CIDAL

## DISINFECTANT - SANITIZER DEODORIZER

EPA Reg. No. 10756-2

### Active Ingredients:

n-Alkyl (60% $C_{14}$ , 30% $C_{16}$ , 5% $C_{12}$ , 5% $C_{18}$ )  
dimethyl benzyl ammonium  
chlorides ..... 5%  
n-Alkyl (68% $C_{12}$ , 32% $C_{14}$ ) dimethyl  
ethylbenzyl ammonium  
chlorides ..... 5%

Inert Ingredients ..... 90%  
100%

For use in HOSPITALS, MORTUARIES, AMBULANCES, CLINICS, NURSING HOMES and INSTITUTIONS, Cidal is a disinfectant, sanitizer, deodorizer and fungicide for use on inanimate hard surfaces where housekeeping will be of prime importance. Its proper use reduces the hazard of cross-infection from environmental surfaces.

CIDAL is a concentrate and should be diluted before using.

## DANGER

**KEEP OUT OF REACH OF CHILDREN**

**SEE RIGHT PANEL FOR FIRST AID  
STATEMENT AND OTHER PRECAUTIONS**

Net Contents 1 Pint  
(16 Fl. Ounces)



**PIERCE CHEMICALS/MORTICIANS SUPPLY**  
409 N. Zang Dallas, Texas 75208

### USE DIRECTIONS

CIDAL will provide the assurance needed in operating rooms, nurseries, autopsy and preparation rooms, restrooms, classrooms, corridors, locker rooms, shower stalls and kitchens. CIDAL may be used for disinfecting floors, walls, sink tops, previously cleaned bed pans, garbage pails, autopsy tables, telephones, etc. For terminal disinfection, use CIDAL at a use dilution of 3½ Oz. per 5 Gal. of water. Apply solution with mop or cloth. Thoroughly wet all surfaces. At this dilution, CIDAL is effective against Staphylococcus Aureus, Salmonella Choleraeausis and Pseudomonas Aeruginosa PRD-10. A dilution of 8 Oz. per 5 Gal. of water is required for fungicidal activity against Trichophyton Interdigitale (a pathogenic fungus causing Athlete's Foot)

Rinse empty container thoroughly with water and discard.

## DANGER

**KEEP OUT OF REACH OF CHILDREN**

Corrosive • Causes eye damage and skin irritation. Do not get in eyes or on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

## FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before re-use.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician immediately.

## NOTE TO PHYSICIAN:

Probably mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

ACCEPTED