

Net Weight 50 Pounds

ARIZONA
INSECTICIDAL MINERAL NUMBER - 35

For Continuous Self-Feeding of Cattle During the Summer for Control of Hornflies and Cattle Grubs; for Mixing in Feed to Provide a 7-day Feeding Program for Grub Control and as an Aid in the Reduction of Cattle Lice, or a 14-day Feeding Program for Grub Control; for Beef Cattle and Non-lactating Dairy Animals Only.

Follow feeding directions carefully.

Pesticidally Active Ingredient:

Ronnel [0,0-dimethyl 0-(2,4,5-trichlorophenyl) phosphorothioate] ---- 6.0%

Pesticidally Inert Ingredients -----94.0%

GUARANTEED ANALYSIS

Calcium (Ca), not less than----- 9.0%
Calcium (Ca), not more than----- 10.8%
Phosphorus (P), not less than----- 9.0%
Salt (NaCl), not less than----- 32.0%
Salt (NaCl), not more than----- 37.0%
Iodine (I), not less than----- 0.00%

INGREDIENTS

Salt, Dicalcium Phosphate, Calcium Carbonate, Propylene Glycol, Magnesium Oxide, Sodium Citrate, Vitamin A Palmitate, Disodium Ethylene Diamine Tetraacetate, Ferrous Sulfate, Manganese Sulfate, Ethylene Diamine Dihydrochloride, Zinc Sulfate, Copper Sulfate, Sodium Sulfate, Cobalt Sulfate, Fumaric Acid, Lactose, Sodium Gluconate, Glycine, L-Lysine.

WARNINGS

Use as sole source of Ronnel.

Dairy cows should be treated early in their dry period but only if the dry period occurs between the end of the adult heel fly season and before grubs reach the back.

Do not feed to dairy animals that will produce milk for human consumption for 10 days prior to freshening. If dairy cows or heifers freshen during feeding program, or if insecticidal mineral has not been withdrawn the required 10 days prior to freshening, milk must not be used for food for 10 days after the last feeding of this product.

Withdraw 10 days prior to slaughter.

Do not feed to sick animals or animals under stress, such as those just shipped, dehorned, castrated, or weaned within the last 3 weeks.

Ronnel is a cholinesterase inhibitor. Do not feed this product to animals simultaneously or within a few days prior to or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals. Consult a veterinarian at the first sign of adverse reaction. Atropine is an antidote.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Wash hands with soap and water after handling. This material contains a cholinesterase inhibitor. Treat as moderately toxic. Atropine is an antidote. This product is toxic to fish, birds, and aquatic life. Do not contaminate water by cleaning of equipment or disposal of wastes. Use this product only as specified on the label. Do not mix with other pesticides. Store in a cool, dry place. Burying in a hole 6 inches deep away from water supplies.

EPA Reg. No. 101-101-01
Licensed Under Federal Insecticide, Fungicide, and Rodenticide Act
© 1974

101-101-01

DIRECTIONS FOR CONTINUOUS SELF-FEEDING
ARIZONA
INSECTICIDAL MINERAL NUMBER - 35

WHEN TO FEED: For hornfly control on pastured cattle start feeding Arizona Insecticidal Mineral Number 35 in the spring before flies appear. Continue feeding all summer and into the fall until cold weather restricts fly development. For grub control, feed continuously for a minimum of 75 days.

HOW MUCH TO FEED: Recommended consumption of Arizona Insecticidal Mineral Number 35 containing 6% Ronnel is 0.2 lb. per 100 lb. of body weight per head per month.

HOW TO FEED: Feed Arizona Insecticidal Mineral Number 35 continuously free-choice to cattle in covered mineral-feeder troughs or boxes. Provide one mineral feeder box or feeding space for each 15 to 20 head of cattle. Locate mineral-feeder boxes and troughs near watering areas. To encourage uniform and normal consumption, put out only a 7-day supply at one time, and keep feeder boxes and troughs clean. For adequate insect control, each animal must eat its share of mineral. Small, late calves may require an additional 7- or 14-day feeding program after weaning for grub control.

WARNING

During continuous feeding program, do not allow cattle to run out of Arizona Insecticidal Mineral Number 35 and do not feed any additional salt or mineral.

DIRECTIONS FOR 7-DAY OR 14-DAY
FEEDING PROGRAMS

Mix Arizona Insecticidal Mineral Number 35 with ground grain for top dressing, or mix with the complete ration to make an insecticidal feed. Feed continuously for 7 or 14 days and be certain that insecticidal feed is evenly distributed on, or mixed into, the total ration. Provide a total of 0.2 lb. of Arizona Insecticidal Mineral Number 35 for each 100 lb. body weight (to be fed over the 7- or 14-day feeding period).

NOTE: Insecticidal feed made from Arizona Insecticidal Mineral Number 35 should be used within 4 weeks of mixing.

WHEN TO FEED: After end of adult heel fly season and before grubs reach the back. Ronnel is not effective for grub control during the later stages of life cycle, when grubs are in the backs of cattle. Reduction of cattle lice will be obtained when fed as directed for the 7-day program.

7-DAY TREATMENT

WHEN TO FEED: Southern area cattle: July through September.
Northern area cattle: August through October.

HOW TO FEED: Mix thoroughly 2 lb. of Arizona Insecticidal Mineral Number 35 containing 6% Ronnel with 10 lb. of feed to produce an insecticidal feed. Prepare only enough for the 7-day feeding program. Feed 0.2 lb. of insecticidal feed per 100 lb. body weight daily for 7 continuous days.

14-DAY TREATMENT

WHEN TO FEED: Southern area cattle: July through September.
Northern area cattle: August through [unclear]

HOW TO FEED: Mix thoroughly 1 lb. of Insecticidal Mineral Number 35 containing 6% Ronnel with 20 lb. of feed to produce an insecticidal feed. Prepare only enough for the 14-day feeding program. Feed 0.2 lb. of insecticidal feed per 100 lb. body weight daily for 14 continuous days.