

**Animal Guard
DIRECTIONS FOR USE**

For Face Flies: Remove mucous secretions and saliva deposits from face of animal. Hold dispenser 18 to 24 inches from animal. Spray face and other body regions where flies gather for 3-4 seconds per animal. Repeat daily, or as needed.

For Horse Flies, Horn Flies, Stable Flies, Deer Flies, Houseflies, Mosquitoes, Gnats: Hold container 18 to 24 inches from animal. Apply lightly to coat with a sweeping motion. Apply to back, belly, ear, or leg regions where the specific attacking insect feeds. Do not spray entire animal. Spray young animals sparingly. Do not apply to back of horses prior to sanding. Repeat daily, or as needed, for protection.

ACTIVE INGREDIENTS:

Apyrethrin.....	.20Z
Piperonyl butoxide, technical.....	.40Z
M-cetyl bicyclohexane dicarbamate.....	.67Z
801-n-propyl isocinchomeronate.....	1.33Z
ADJUVANT INGREDIENTS.....	97.40Z
Total	100.00Z

*Consists of .32Z (butylcarbitol) (6-propylpiperonyl) ether and .08Z related compounds.

**MCA Repellent 326, U.S. Pat No. 2,757,120

***contains petroleum distillate

CAUTION

HAZARDOUS IF SWALLOWED. Do not contaminate feeds, water or mineral supplements with this spray. Do not contaminate milk or milk handling equipment with this spray. Wash udder and teats before milking. Avoid overspraying which will result in wetting animals hide.

PHYSICAL OR CHEMICAL HAZARDS

EXTREMELY FLAMMABLE. Contents under pressure. Keep away from fire, sparks and heated surfaces. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

PM 17

10806-17

12/14/90

1/1

**ANIMAL
GUARD**



EPA Est. 10806-NJ-1
EPA Reg. No. 10806-17

**FOR TEMPORARY PROTECTION OF DAIRY
CATTLE, BEEF CATTLE, AND HORSES FROM
ATTACK BY HORSEFLIES — HORN FLIES —
STABLE FLIES — DEER FLIES — HOUSEFLIES
— MOSQUITOES — AND GNATS.**

**CAUTION:
EXTREMELY FLAMMABLE**
See other cautions on back panel.
KEEP OUT OF REACH OF CHILDREN.

BEST AVAILABLE COPY

