COUTSON AFRICAN VIOLET SERAY

For Listed Insects on Ornamental Plants

ACTIVE INGREDIENTS: Tetramethrin [(1-Cyclohexene-1,2-dicarboximido)methyl 2,2-dimethyl-3-(2-methylpropenyl)cyclopropanecarboxylateJ..... .0286% *3-phenoxybenzyl d-cis and trans**2,2-dimethyl-3-(2-methylpropenyl) .0547% *Other isomers..... .0025% INERT INGREDIENTS..... 99.9142% 100.0000%

*d-(cis, trans)phenothrin **cis/trans isomer ratio: max .25% (+ or -) cis min .75% (+ or -) trans

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

ON PLANTS: Kills Spider Mites, Aphids, Mealybugs, Japanese Beetles and others listed on back panel.

Net Weight: 6 Ounces

Manufactured by: Contact Industries 641 Dowd Ave. Elizabeth, N.S. 07201 10806-57

EPA Est. /0806 -NJ-/

EPA Reg. No./0806-57

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Avoid breathing vapors. Wash hands with soap and water after use.

Do not use in commercial food processing or preparation areas. In the home, all food processing surfaces, exposed food and utensils, should be covered during treatment, or thoroughly washed before use. Do not use on edible plants.

Cover fish aquariums before spraying.

PHYSICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame, Do not puncture or incinerate container. Exposure to temperatures above 130 F may gause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Store in a cool area away from heat or open flame. Storage:

Disposal: Do not reuse empty container. Wrap container and put in trash collection.

SHAKE BEFORE USE. HOLD CONTAINER UPRIGHT.

OR USE ON ORNAMENTAL PLANTS, INDOORS AND OUTDOORS:

Pepperonia, Spirea, Philadendron, African Violet, Euonymus, Seedum, Ferns, Crabapple Rose, Wisteria.

When sprayed directly on insects, this product kills sucking insects: Aphids, Spider Mites, Leaf Hoppers, Whiteflies and Mealybugs. Kills chewing insects: Japanese Beetles and Southern Army Worms.

FOR INDOOR PLANTS:

Spray both upper and lower surface of leaves from a distance of 12-14 inches until visibly wet. Repeat as needed.

FOR OUTDOOR PLANTS:

For best results spray in the evening or on a cloudy day when air is still. Avoid using in hot, bright sunshine. Use slow sweeping motion, holding the dispenser approximately one foot from foliage. Spray both upper and lower leaf surfaces until visibly wet. Repeat as needed.