MISTY TIME DISPENSER Concentrated Aerosof baser Killer contains a highly effective insecticide formulation speed fly designed for use in automatic mist dispensing units around food handling establishments, Food Processing Establishments, test rooms, dining rooms and restaurants where flies, gnats, flying moths, and mosquitoes are an annoyance.

ACTIVE INGREDIENTS:

Pyrethrins								
Piperonyl Butoxide Technical*					,			2.00%
N-octyl Bicycloheptene dicarpox	in	ni	de			,		3.34%
Petroleum Hydrocarbons								13.66%
INERT INGREDIENTS								80.00%

 Equivalent to 1.6% (butylcarbityl) (6-propylpiperonyl) ether and 4% related compounds.

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator.

CAUTION: Do not use in nurseries or rooms where infants, ill or aged persons are confined. Do not install directly over any food handling or dispensing areas. Large drafts of forced ventilation may reduce or eliminate the insecticidal efficacy of these installations.

MISTY MIZIER INSECTICIDE

Mfg. By AERO MIST, IMC. Industrial Pk. Dr.

ndustir,al Pk. Dr. Marietta, Ga. **30062**

Net Wt. 12 oz.

E.P.A. No. 10307-27

CAUTION

Keep out of reach of children. See side panel for additional cautions.

of USE IN AUTOMIZER: One Automizer unit of ould be inshall de for each 6,000 cubic feet of, space to be treated. The Automizer activates the 100mg valve once every 15 minutes; είν του hours of operation, enough insecticide is suspended in the air to kill a substantial proportion of the flies, mosquitoes and gnation resent. Usually the effectiveness will be noticeable within can hour after the Automizer is activated. If less than 24 hour consist renormed, the Automizer should be set to start one hour. before control is desired and run one hour after control is no ; longer needed. Keep in mind that such factors as air flow rate," size of space treated, humidity and room temperatures all effect insigt activity and control. Therefore, one unit per 6,000 cubic. feet should be considered as an average need with adjustments made depending on specific conditions at place of usage. As with any method of insect control, best results are achieved when sanitation and other positive insect-prevention practices are followed.

Thi, 12 oz. can of insect killer contains approximately 3,300 "shots" sufficient to sustain operation for a total of 825 hours (34 days on around the clock basis) or 68 days on a 12 hour basis. When used on a 24 hour basis, it is suggested that the container be changed once a month (once every 2 months on a 12 hour basis.)

ACCEPTED

WAR 2 7 1973

UNDER THE FACULAL INSECTICIDE FUNCTION PORCHAM NUMBER AND NUMBER OF THE PORCHAMAN NUMBER OF THE PORCHA