

WISLEY

INDUSTRIAL AQUEOUS EMULSIFIABLE CONCENTRATE

An Insecticide Spray Concentrate

For use in Food Processing & Storage, Mushroom
Houses, Hospitals, Hotels, Motels, Restaurants,
other Institutions and in Residences

Authorized for use in all departments of official establish-
ments operating under the meat, poultry, shell egg grading
and egg products inspection programs.

Provides Rapid Kill of Insects Without Leaving
Objectionable Residues

Food additive tolerances have been established for the
insecticide components in this product. When this product is
used as directed, allowable tolerance levels will not
be exceeded

ACTIVE INGREDIENTS:	
Pyrethrins	3.0%
*Piperonyl Butoxide Technical	30.0%
Petroleum Distillate	12.0%
INERT INGREDIENTS:	
	55.0%
	100.0%

*Equivalent to 24% (butylcarbityl(6-propylpiperonyl) ether and
6% related compounds

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See side panel for additional precautions.

ONLY FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS

EPA Reg. No. 10807-RER EPA Est. No. 10807-GA-1/10807-TX-1

Manufactured By

AMREP, INC. • 990 Industrial Park Dr • Marietta, GA 30062

NET CONTENTS GALLON(S)

To control carpet beetles, spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

To control fleas, and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

As a Space Spray: Dilute at rate of 1 part concentrate to 59 parts water (2 fluid ounces per gallon). Apply with mechanical or compressed air equipment (non-thermal) adjusted to deliver a fine mist. Close doors and windows and shut off ventilating systems. For rapid control of adult house flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths: Direct spray at an upward angle distributing it uniformly through the area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for at least 10 minutes. Ventilate and sweep up dead insects before reoccupying.

For rapid kill of exposed or accessible stages of other insects named on this label, dilute 1 part concentrate in 5-29 parts water (4 to 25 ounces per gallon). Apply using conventional mechanical or compressed air equipment (non-thermal), following directions for space spraying. Apply the above dilutions at a rate of from 1 to 10 ounces per cubic feet of space. When used in meat and poultry processing plants, a dilution of no less than 1 to 5 is acceptable and it must be dispersed from equipment adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns).

STORAGE and DISPOSAL

Prohibitions: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.
Pesticide Disposal: Pesticide spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State and local procedures.
Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in sanitary landfill or by other approved State and local procedures.