

DIRECTIONS FOR USE: It is a violation of Federal law to use the product in a manner inconsistent with its labeling. **SHAKE BEFORE USING AND AT INTERVALS DURING PROLONGED USE.** Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range. **HOUSE FLIES, MOSQUITOES, GNATS, WASPS AND SMALL FLYING MOTHS (MILLERS):** Close all doors and windows and spray room at the rate of 2 seconds per 1000 cubic feet. Direct spray to all parts of the room especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Repeat as necessary. **COCKROACHES, WATERBUGS, SILVERFISH, CRICKETS, SPIDERS, CENTIPEDES AND SOW BUGS:** Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary. **ANTS:** Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary. To kill **FLEAS** and **BROWN DOG TICKS** on dogs, cats, and premises: Apply directly to animals along back, under forelegs and on other areas except genitalia. Part hair to contact skin and insure good cover. Hold nozzle about one foot away while spraying. spray the animal's bedding and other resting areas until the surface is slightly moist. Also spray the entire inside surface of dog houses and kennels, thoroughly spraying floors, cracks, crevices and bedding. Hit as many insects directly as possible. Repeat as necessary. **CARPET BEETLES:** Spray directly on beetles, baseboards, floors, shelves, and carpets. Repeat as necessary. **BEDBUGS:** Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Allow treated fabric to dry thoroughly before using. Repeat treatment as necessary. **IN ANIMAL QUARTERS AND MILK ROOMS:** For rapid control of **HOUSE FLIES, MOSQUITOES, GNATS, WASPS AND SMALL FLYING MOTHS:** Close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed. **TO TREAT ANIMALS:** To protect animals from attacks of **STABLE FLIES, HORSE FLIES, DEER FLIES, HOUSE FLIES, HORN FLIES, MOSQUITOES AND GNATS:** Spray about 3 seconds to each side being careful to spray back withers and forelegs thoroughly. To protect from **FACE FLIES:** Spray the face and head, but do not spray into the eyes. Repeat treatment as necessary when flies are troublesome. **EXPOSED STAGES OF INDIAN MEAL MOTHS AND ANGIOMOIS GRAIN MOTHS:** Close doors and windows. Direct spray into all parts of room, especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for that purpose. **EXPOSED STAGES OF FLOUR BEETLES, WEEVILS AND GRAIN MITES:** Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary. Infested stored products in the home should be disposed of. **CONTROL GARDEN INSECTS ON ORNAMENTAL AND VEGETABLE PLANTS:** Spray with slow sweeping motion, not closer than 18 inches from plants. Avoid wetting plants. Spray plants only when air is still. For single plants, spray in short bursts. Repeat as necessary for control. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90° F. Contact insects directly whenever possible. Repeat application at 7 to 10 day intervals or after heavy rain. Acceptable for Use to Day of Harvest on Edible Plants. Use on Ornamentals (Roses, Asters, Dahlias, African Violets, Azaleas, Euonymus, Hydrangeas, Junipers, Lilacs) to control **APHIDS, ROSE CHAFERS, JAPANESE BEETLES, ROSE SLUGS, TENT CATERPILLARS, SCALES (including EUONYMUS and JUNIPER SCALES), LEAFHOPPERS, TARNISH PLANT BUGS, LACE BUGS (including AZALEA LACE BUGS, THRIPS (exposed), MEALY BUGS (including CITRUS MEALYBUGS), MITES (including RED SPIDER, SPIDER and BROAD MITES), PLANT BUGS, WHITEFLIES, LEAF MINERS, HYDRANGEA LEAF TIERS, BAGWORMS and LEAF ROLLERS.** Use on Vegetables (Peppers, Eggplants, Potatoes, Cucumbers, Squash, Melons, Beans, Sweet Corn) and Tomatoes to control **FLEA BEETLES, HORNWORMS, STINK BUGS, COLORADO POTATO BEETLES, APHIDS (including GREEN PEACH APHIDS), LEAFHOPPERS, BLISTER BEETLES, STRIPED and SPOTTED CUCUMBER BEETLES, MEXICAN BEAN BEETLES, WHITEFLIES, 12-SPOTTED BEETLES and JAPANESE BEETLES.**

STORAGE AND DISPOSAL: Store in a cool dry area away from heat or open flame. Do not transport or store below 32°F. Replace cap, wrap container in several layers of newspaper and discard in trash.
PRECAUTIONARY STATEMENT: HAZARDS TO HUMANS AND DOMESTIC ANIMALS:
CAUTION: Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in food processing plants, wash all equipment, benches, shelving, etc. where exposed food will be handled, with an effective cleaning compound followed by a potable water rinse before reuse. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Remove pets, birds, and cover fish aquariums before spraying. In hospitals, remove patients prior to treatment. Ventilate rooms for 2 hours before returning patients.
STATEMENT OF PRACTICAL TREATMENT: IF SWALLOWED: Contact a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching finger to back of throat. Never give anything by mouth to an unconscious or convulsing person. **IF INHALED:** Remove affected person to fresh air. Apply artificial respiration if indicated. **IF IN EYES OR ON SKIN:** Flush affected areas with plenty of water. Contact a physician if irritation persists.
ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water. Do not contaminate water by disposal of wastes or cleaning of equipment.
PHYSICAL OR CHEMICAL HAZARDS: Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

EPA Reg No 10807-156 EPA Est. No 10807-GA-1

11:33



Manufactured by:
 Amrep Inc.
 888 Industrial Park Drive
 Marietta, GA 30062

Amrep 5003

- For Use In: Schools, Hospitals, Institutions, Supermarkets, Dairies, Hotels, Restaurants, Drive-Ins, Motels, Drug Stores, Food Plants, Milk Rooms and Animal Quarters.
- Provides Rapid Control of: House Flies, Mosquitoes, Small Flying Moths (Millers), Gnats, Wasps and accessible stages of Cockroaches, Ants, Waterbugs, Silverfish, Flour Beetles, Weevils, Grain Mites and Spiders.
- Kills and Repels: Stable Flies, Horse Flies, Face Flies, Deer Flies, House Flies, Mosquitoes and Gnats.

ACCEPTED
 2 JUN 1994
 for the Federal Insecticide, Fungicide, and Rodenticide Act
 registered for the pesticide
 10807-156

ACTIVE INGREDIENTS:
 Pyrethrins 0%
 *Piperonyl Butoxide, Technical 5%
INERT INGREDIENTS: 94%
TOTAL: 100%
 *Equivalent to 4% (butylcarbityl)(6-propylpiperonyl)ethers and 1% related compounds

KEEP OUT OF REACH OF CHILDREN.
CAUTION:
 See back panel for additional precautions.
NET WT. 16 OZ. (454 GRAMS)
 (Equivalent to 19.7 Fluid Ounces)

(11-05-93 copy to submit to EPA)

BEST AVAILABLE COPY