10807-188

10/03/2003



## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

OCT : 3 2003

ŧ

ł

Ms. Susan Watson Regulatory Manager Amrep, Inc. 990 Industrial Park Drive Marietta, GA 30062

Subject: Response to Agency's August 12, 2003 Letter Concerning Label Amendment to Update First Aid Statements (PR Notice 2001-1) and Disposal Instructions (PR Notice 2001-6)

> EPA Reg. No. 10807-188 Misty Industrial Insect Killer Your Submission Dated August 25, 2003

Dear Ms. Watson:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comments listed below. Two (2) copies of the finished labeling must be submitted prior to releasing the product for shipment. A stamped copy of the label is enclosed for your records.

- 1. Revise the statement "Do not use on puppies under 4 weeks" to read: " Do not use on puppies under 12 weeks". Domestic animal safety data are required to support use on younger animals.
- 2. Considering label space restrictions we recommend you add the word "medicated" to the first statement beginning "Consult a veterinarian before using this product on debilitated...etc.", allowing deletion of the last two sentences regarding medication.
- 3. Since there are more than just pet uses on the label we recommend you move the statement "Read entire label before each use" to the front panel.

If you have any questions regarding this action, please contact Olga Odiott of my team at (703) 308-9369.

Sincere h George TV LaRocca

2/3

Product Manager (13) Insecticide Branch Registration Division (7505C)

Enclosure

2

(

PRECAUTION: Harmiu // Inhaded. Avoid breathing vapor or spray mist. RAIZARIOS TO HEIMANS AND DOMESTIC ANNIALS CAUTION: Harmiu // Inhaded. Avoid breathing vapor or spray mist. Remove contaminated clothing and wesh before reuse. FIRST AND STATEMENT: If INMALED: Move person to treat air. If person is not breathing, call 811 or an ambulance, then give artificial respiration, preferably mouth-to-mouth Harm the product consister or bedue with very dorive fragment article.	
There are product contraries or allow with you write towing a posteri control rearies or uponts for interminent ENTROMMENTA. NAZARDS: This product is toxic to lish. Do not apply directly to any body of water you not contraminate water when disposing of equipment washwaters. PHYSIGCA OR THERMOLEL HAZARDS: Contration of the prostore to not use of the need of the production of purplet of equipment estimates above 130°F may cause bursting. Do not use this product in or on electrical equipment due to the possibility of shorts hazard.	
DIRECTIONS FOR USE: IT IS A VARIATION OF FORMER FOR USE THIS PLOUGH IN A THEMAN STATUSSIENT WIT IS AUTOMING TO STODAGE & DISPOSALL: STOPAGE: Store in a cool, dry area. Do not itansport or store below 32FF. CONTAINER DISPOSAL: DO Not Puncture of incidential IF EMPTY. Place in treash or object for recycling if arealable. IF PARTIX FILLED: Call your local solid waste agency or 1- 800.CLEMUP for disposal instructions.	900 Industrial Part Drive Marieta, GA 20062
Do not use in foordiead areas of foordiead handling establishments, restaurants or other areas where loordiead areas of foordiead areas of processed. Do not use in serving areas are areas where areas where hows are served processed. Do not use in serving areas areas where hows may be propared or had. Nontoordinaride areas where propared to not are served such an excit of a served areas where hows where hows may be propared or had. Nontoordinaride areas are areas are areas areas areas areas areas areas areas areas areas of the formation of the areas are	
SHAKE WELL BEFORE USING. Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches. Avoid containination of lead and loodstaffs. Remove pets and birds and cover lish aquanium before spraying. Vacate room after treatment and vertiliste before reoccuping. Do not allow children or pets to contract treated areas until surfaces are dry. NOOORS: Cleater Firet, Nonevilles, Fruit Rive, Moorgulica, Greats, Mapp, Hornats, Bees, Black Firet, Skipper Files, Blow Files and Small Flying Moths: Close all doors and encloses and direct spray upward into center of room with a slow server 5-10 seconds Small Flying Moths: Close all doors and encloses and direct spray upward into center of room with a slow server motion. Spray 5-10 seconds	
pri everage occur, more grower or occi. Livera spay to all parts of the norm estade whore and offen process whore affect the seven affect the seven and affect the seven affect of the sev	Industrial Inse
Spray rate, nests and points of entry. Spray on entry when possible. Repeat as necessary indian liked likethe, Angeumoté Grein likether. Close doors and windres, direct spray into all petrs of noom especially around strond lood product containent. In paties and duriened area. Spray 7 to 19 asconds for each 1000 outois level of spcar. Noving strond strond and liketh area data for Belles and duriened areas. Spray 7 to application as necessary. Exponent dataper of contraved from Beather, Charge area charder to far Beather, Graner application as necessary. Exponent and a novie prevent Specifies (Specifie Beather, Drag Spare Beather, Spray and and and and and an examination and an examination and and and and and and and and and an	
or containers or other areas where these insects tend to construct as many insects as possible. Spray 30 seconds for each 100 space tend of surface. Keep area closed for 45 minutes to 1 hour after treatment. Spece Spray: 30 seconds per 1000 cu. R. of space, moving towerd the door team nom and keep area closed for 15 minutes holowing treatment. Spece Aprex application weep or an exact moving towerd the door team nom and keep area closed for 15 minutes holowing treatment, they are possible to a for an and keep area to a for a spece. The door team of team of the door team of the door team of team of team of the door team of the door team of team of team of team of the door team of the door team of team o	• Controls Industrial Insects • Socks Out Rugs Where -
process strong or graphone up an exprove proving treatment on eacy provide verse provide and and an another process the process period of the providence and other processing the providence of	624
Davis motoughy, and as not several neurs in sumpril, is possible, Apply spiry moreary to ease, correct, and the market are correcting and the second several provided and the second several provided and the second several provided and the several provided and t several provided and the several p	
and vertiated before neoccupring. Repeat treatment after 7 days or as needed. Pleas & Tichs (Adelit and Larvas); Remove and Shooudyby clean or destroy pet bedding. Syray per supplies quertes, bad, entire Skor and Skor on Skor age with special attention to creates and creates, areand baseboards, and window and door iname. Indentic upholatend familien and matterease range date be treated with a poticit. A test application should be made to upholatent or depoint fairlies in all inconductions with special attention to create and creates are application should be made to upholatent or depoint fairlies in all inconductors attentions range date be treated with the product. A test application should be made to upholatent or depoint fairlies in all inconductors takes to the source of the attention be attended and the attended state attended	
head, strote against the lay of the heir, spraying the parted for it short bursts, diesdly behind the heard is heare panetrating the Cost. Dampon the two to be ust. MOTE: REJADENTRE LABEL BEFORE EACH USE. USE ONLY ON DOGS. Reappy weakly increasary to control structuron. Do Not Use On Parpise Under A Weeks. Consult a wateriation before using this product on deblackate. Jobel, parter	Scorpions, Silverfick, Soudnugs, Spiders, Ticks, Wasps, 4949, Other Peats Listed on the Label D.V. D.V.S.
of verse. If styre continue, consult a vestmentian immediately. Cestuin medications can interest with perticities. Consult a vestmentian before using on medicated animals. To tail Libe and Levere agger Spery in an incomplications area to test for possible staining or discritionality. Imspect again after christige then proceed to styre write verse how to be treated. Hold container work with write an exclusion area to test for possible staining or discritication. Imspect again discriting then proceed to styre write verse how to be treated. Hold container work with promote and outs of write verse where waters and discritications of 8 to 10 stribus. Some west strane boot for the seconds. Some other than an examption and and and outs of hordinon hordinon and stray from a	EPA Rag No. 10607-188 EPA Est. No. 10607-6A-1 EP H
kumbur the carnot be either laundwerd or dry channed. Alther all strayed articles to dry throwughly before use. IN ANBULL OUARTERS AND NEXT ROOMS: For rapid combrol of housefflee, likoequiboee, Gmate, Wanne, Bees and Samil Frying Biother: Close windows and doors and spray all the rate of 2 seconds per 1000 cubic feet of space. For best results, keep and choose differ spray de	AcTINE MEMBERINE Productos Technical
ans and verifiele dist treatment is completed. To TREAT LIVESTOCK: To protect built callet, dairy callet, goats, aboue, hoge and horses from attacht of Sabib Flins, Hone Flins, Done Flins, Foco Flins, Nouvelles, Hon Flins, libogallons, and Gmits: Spay dool 3 second on each being carried to spay back, where and horselps. To protect from Flins, Spay the kine and head, but on tapoy isto area. Spanse treatment who first are anti-horsens from ships. The anti-horselphone, and frank but do not spay isto area. Spanse treatment who first are anti-horsens and some Stream. For a spanse first schement 10 COMITED	"Permetrin More Construction (2005) A mart more construction (
BLOOD SUCKING LLCE: Apply to he Meased avear of the arintal, using a still breach to get the spore for the base of the fault. Repeat every 3 weeks # required. TO CONTROL POULTRY LUCE: Sporey roots, wells and needs or cages thoroughly. This should be followed by sporying over the birds with a factor and the CONTROL of SEDBUGS AND INTES IN POULTRY MOUSES. Spory creates of noost poles, creates in wells, and creats in needs above the burdent and mast heat.	"Equivalent to min. 0.1% burk/saffagi (5 propytoecony) either and 0.1% related compounds. "(5 phenoxypheny) melly (++) disfrants 3-(2.2 dichthroathany) 2.2 dimethylichproperecationytate. Cisthrans ratio: min. 30% (++) dis and mar. 65% (+4) hans.
OUTDONG: Point sparsy hour lace. Hold can on slight downwed angle. For best results, sparsy when at is calm. Allow a law minutes for product to take effect. Spray with wind it breaze is blowing FLYING INSECTS: Files, Mongathoes, Ganat, Sanall Phyling Moths, Hormeta, Waage and Yellow Allow Koji of the are area. Direct spray loward busines or goes from at least 40.6 free special advention to breading and Yellow	KEEP OUT OF REACH OF CHILDREN.
arees such as garage cars and compose pres. Hepen, as necessary, must bencer and want heart: Spray must one cay on traces and may mark mass from approximately 2 feet aney. Considering the mark into nog Applications should be made in the evening with an a frast Spray into Mading and Dreeding places contacting as many insects as possible. CAMILING MISECTS: Ante, Controches, Mode	CAUTION:
possible. Also spray legs of tables and chairs to repellants. Spray and hills for best control.	See back panel for additional precautionary statements. NET WT. 16 OZ. (1 LB.)(454 GRAMS)

100 ×

CO\25\80 to as pob.(qC0-80)881-70801 oN geA AGE/coores/1

NOTE: Areas highlighted in blue are per Agency letter dated 08/12/03.