



U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460

NOTICE OF PESTICIDE:

x_ Registration Reregistration 10807-199

EPA Reg. Number:

FEB 2.4 2009

Term of Issuance: Unconditional

Date of Issuance

Name of Pesticide Product:

Misty Dualcide P3 RTU

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Amrep, Inc 990 Industrial Park Drive Marietta, GA 30062

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is <u>unconditionally</u> registered in accordance with FIFRA sec. 3(c)(5), provided that you submit two (2) copies of your final printed label before you release the product for shipment. Please refer to the A-79 enclosure for a further description of final printed labeling. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing amended labeling constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. If you have any questions, please contact Dr. William Sproat of my team at (703) 308-8587.

Signature of Approving Official:

Date:

FEB 2 4 2003

George T. LaRocca

PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION: Wash thoroughly with soap and water after handling^. ^

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high water mark. Do not contaminate water when disposing of equipment washwaters.

STORAGE & DISPOSAL: Do not contaminate water, food or feed by storage or disposal, PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste. PESTICIDE DISPOSAL: Pesticide or finse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility. CONTAINER DISPOSAL: Containers one gallon and amalter: Do not reuse container. Wrap container in several layers of newspaper and facard in trash. Containers larger than one gallon: Metal Containers: Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities. Plastic Containers: Triple rinse or equivalent, then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, of if allowed by state and local authorities, by burning. If burned, stay out of smoke. Glass Containers: Triple rinse or equivalent, then dispose of in a sanitary landfill or by other approved state and local procedures.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE EACH USE PET USE APPLIES TO USE ONLY ON DOGS


Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ormanental gardens or parks, or on golf courses or lawns and grounds. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover lish equaria before space spraying or surface applications. Do not use on humans. Vacate room after treatment and ventilate before reocupying. On not allow children or pets to contact treated areas until surfaces are dry. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing animals. Do not use on dogs under 12 weeks. Sensitivities may occur after using ANY pesticice product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

SHAKE WELL BEFORE USING.

INDOOR APPLICATION:

Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held. In the home, all food processing surface and utensits should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed. Nonfood/feed areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or botting). Vacate treated areas and ventilate before reoccupying. For maximum effectiveness, a combination of spot, surface treatment and fogging is recommended. Do not apply to classrooms while in use. In the home, all food processing surfaces and utensits should be covered during treatment or thoroughly washed before use. Do not apply this product in coccupied patient rooms or in any rooms/areas while occupied by patients, the elderly or infilm. Do not apply to institutions (including libraries, sports facilities, etc...) in the immediate area when occupants are present. Exposed food/feed should be covered or removed.

SURFACE SPRAYING: For 60-Day (Two Months) control Cockroaches, Waterburgs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverlish, Firebrata, Solders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetles, Meal Worms, Grain Mites and Cadelles: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nest and points of entry. To control Carpet Beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and intested areas of shelving. Repeat treatment as needed. To control Bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, molding and floors. Repeat treatment as needed. To control Floor (adult and larvae) and Ticks, including Ticks which may carry Lyme Disease (adults and larvae): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire area of floor and floor covening. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas. To control Clothes Moths: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible.



Dualcide P3 RTU

- For Use in Homes and Non-Food Areas of Food/Feed Processing Establishments, Kennels, Commercial Buildings, Schools, Motels, and Hotels.
- For Control of Fleas, Brown Dog Ticks and Lice on Premises and on Dogs
- For Control of Ticks Which May Cause Lyme Disease on Premises
- Controls Fleas and Ticks on Dogs for 14 Days
- For Use Both Indoors and Outdoors
- Provides 60-day (Two Months) Control of Cockroaches,
 Fleas, and Stored Product Pests

EPA REG. NO. 10807-199 EPA EST, NO. 10807-TX-1

ACCEPTED

FEB 2 4 2003

ACTIVE INGREDIENTS:

"Permethrin: 0.50%

MERT INGREDIENTS†: 99.50% Under the Pederal Insecticide,

TOTAL: 100.00 Fungicide, and Rodenticide Act.

"(3-phenoxyphenyi) methyl (±) cis/trans 3-(2.2-dichloroethenyl)
dimethylcyclopropanecarboxylate. Cis/trans ratio: min. 35% (±) cis and max. 65% (±)

"Gegintered under
EPA Req. No. 10807 - 199

KEEP OUT OF REACH OF CHILDREN.

CAUTION:

B00442

See side panel for additional precautionary statements.

NET CONTENTS 1 GALLON (3.8 LITERS)

Apply spray littlerally to empty chests, or cracks, joints and crevices. After airing, i three feet from fabric to avoid staining. 7 to test for possible staining or discoloration to be treated. Soray from a distance of 8 including mattresses and furniture that articles to dry thoroughly before use. SPACE SPRAYING: Close doors and y machine adjusted to deliver a tine mist or Mosquitoes, Skipper Flies, Waspe, Mediterraan Flour Moths, Indian Me angel, distributing it uniformly throughout the great closed for 2 hours. Venillate it FOGGING: To kill exposed or accessit of Ligation to 20,000 to 40,000 cubic feet closed for 2 hours. Ventilate before reco ON LIVESTOCK: To protect cattle (be Files, Face Files, House Files, Mosq surface of the hair. For face files, soray f Files, Horse Files and Deer Files: App hair thoroughly. Repeat treatment daily control Blood-Suciding Lice: Apply to I soray to the base of the hair. Reneat ever walls and nests or cages thoroughly. Th mist. To control Bedbugs and Mites is walls, and cracks in nests where the be ON DOGS: To control Flees, Ticks and and making sure that the animal's entire spraying, fluff the hair so that the spray wi do not saturate animal. Do not spray inti gloves or mitts when applying product. F USE ON CATS OF KITTENS.

To control Ticks, including Ticks which product as a residual treatment to contro Star Tick (Amblyomme americanum) can Dog Tick (Dermacentor variabilis each areas listed below, apply until surfa 750-1000 square feet). Treat around win under stätnwavs and in crawl spaces wher litter 6 to 10 feet wide around the outside of ticks in outdoor activity areas, treat a ban activity areas, especially where dense ve on roses, dahlias, asters and other ornar Whiteflies, Aphids, Exposed Thrips, S as a multi-purpose soray in small carder prevent spreading in large garden areas asters, azaleas, begonias, camellias, ca woods, English ivy, eunonymus, faschia. marigolds, rhododendrons, roses, rubbei Effective when used as directed again Thrips, Lace Bugs, Japanese Beetles spraying plants, do not operate closer than tender foliage, young plants and new gro Contact insects directly whenever possi oughly treat the entire area. For best resul hour but do not apply when wind speeds ex mosquitoes may rest. Apply at a rate of 2.5 mounted conventional misting equipmen mph to treat a 300 foot swath, calibrate eq. should open water be treated with this pro-

Buyer assumes all risks of use, storage directions given herewith.

frmoc-pend/epa-pend/10807-199/Aventisas of 12/04/02 (areas revised per EPA let