



LCF

NTS:

FOR ANIMALS:

or absorbed through skin or eyes. Avoid breathing spray

handlers and other handlers must wear - Chemical resistant gloves (nitrile) - Chemical resistant eye protection - Chemical resistant headgear - A dust/mist filtering (P100), or a NIOSH approved

Follow the instructions for cleaning/ use detergent and hot water. Discard clothing and other gear heavily contaminated with this product. Use closed systems or enclosures in the Worker Protection Standard (29 CFR 1910.120), the handler PPE in the WPS.

Remove clothing immediately if contaminated. - Users should wash the outside of gloves before changing into clean clothing. - Product should be washed in a separate container from other clothing, chewing gum, using

3. Rinse skin immediately with water and soap. Call a poison control center or doctor for treatment if necessary. Do not induce vomiting. Do not swallow. Do not induce emesis. IF INHALED: Move person to fresh air. If breathing is difficult, call a poison control center or doctor

for further treatment advice. IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container label with you when calling a poison control center or doctor or going for treatment.

ENVIRONMENTAL HAZARDS (TERRESTRIAL AND AQUATIC USES): This pesticide is toxic to aquatic invertebrates. Do not apply to water except as specified on this label. For **Terrestrial Uses:** Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. For **Aquatic Uses:** Do not apply directly to water except as specified on this label.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Do not stack more than four drums high. Leaking or damaged drums should be placed in overpack drums for disposal. Spills should be absorbed in sawdust or sand and disposed of in a sanitary landfill. Keep container closed when not in use. Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store at temperatures above 32°F. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

GENERAL PRECAUTIONS AND RESTRICTIONS: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

ENTRY RESTRICTIONS: For 4 days following applications to non-crop areas (other than aquatic or residential sites), do not allow employees to have contact with the treated plants, except for contact with their footwear. Do not allow people or pets to touch treated plants until the spray has dried. - Do not use the treated water for animal consumption, spraying, irrigation, or domestic purposes for at least 5 days after treatment. - Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption. Do not contaminate water by cleaning or equipment or disposal of wastes. - Do not contaminate feed, foodstuffs or drinking water. - Do not store or transport with feed or food.

APPLICATION INSTRUCTIONS (Site Specific):

NON-CROP OR NON-PLANTED AREA USES: To kill above ground weed growth around buildings and other installations, ornamental gardens, walkways, patios, beneath greenhouse benches, fence lines, golf courses, parkways and driveways for control of undesirable broadleaved and grassy weeds - 1 gal. per 20 gal. of water (6 1/2 oz. per gal. water). Apply for full coverage and thorough weed contact. Apply preferably to young growth and up to seeding of weeds. Repeat applications necessary to maintain control, particularly on grasses and established weeds. Avoid spray contact with foliage or food crops or ornamental plants. Do not water treated areas for 24 to 36 hours after application.

AQUATIC USE DIRECTIONS: For application only to still water (i.e., ponds, lakes, and drainage ditches) where there is little or no outflow of water. Do not use the treated water for animal consumption, spraying, irrigation, or domestic purposes for 5 days after treatment. Treatment of dense weed areas can result in oxygen loss from decomposition of dead weeds. This loss may cause fish suffocation. Therefore, treat only 1/3 to 1/2 of the water body area at one time, especially if dense areas of weeds and/or algae exist, and wait 14 days between treatments. Consult your State Fish and Game Agencies before applying this product to public waters. DO NOT APPLY TO MUDDY WATER. Avoid creating muddy water during application. Do not apply under conditions of high wind and wave action. Do not apply where plants are covered with mud deposits.

WEED SPECIES	GALS./SURFACE ACRE
SUBMERGED WEEDS (infesting Still Ponds, Lakes, Ditches, or Portions Thereof):	
Bladderwort (<i>Utricularia</i> spp.)	10 - 20
Coontail (<i>Ceratophyllum demersum</i>)	20
Elodea (<i>Elodea</i> spp.)	20
Naiad (<i>Najas</i> spp.)	10
Pondweeds (<i>Potamogeton</i> spp.)*	20
* Except Richardson Pondweed (<i>R. richardsonii</i>) in Minnesota	
Watermilfoil (<i>Myriophyllum</i> spp.)	10 - 20

APPLICATION FOR SUBMERGED WEEDS: In early season where submerged weed growth has not reached the surface apply this product by pouring directly from the container into the water while moving slowly over the water surface in a boat. Distribute evenly over infested areas in strips 40 feet apart. In mixed weed populations, use the high rate of application as indicated by weeds present.

In late season or where submerged weed growth has reached the water surface throughout the area to be treated apply this product by pouring as above or injecting a diluted (10-20 to 1) solution below the water surface in strips no more than 20 feet apart. Use the high rate of application as indicated by the weeds present. Where algae are present along with the submerged weeds, pre-treatment with Copper Sulfate at recommended rates is advised for best results with this product.

FLOATING WEEDS	GALS./SURFACE ACRE
Pennywort (<i>Hydrocotyle</i> spp.)	5 - 7 1/2
Salvinia (<i>Salvinia rotundifolia</i>)	5 - 7 1/2
Waterhyacinth (<i>Elodea crassipes</i>)	5 - 7 1/2
Waterlettuce (<i>Pistia stratiotes</i>)	5 - 7 1/2

APPLICATION FOR FLOATING WEEDS: 150 to 200 gallons of water plus a non-ionic surfactant* should be applied as an overall spray for control of the above weeds. The high rates of this product should be used for late season application.

*Add a Nonionic Surfactant containing 75% or greater surface active agent at the rate of 1/2 to 4 pints per 100 gallons of finished product.

Duckweed (*Lemna* spp.) 10
APPLICATION FOR DUCKWEED: For control, apply this product as an all-over spray in 50 to 150 gallons of water per acre. Care should be taken to thoroughly cover all plants on water and on damp marginal areas. Reinfestation of duckweed occurs readily from untreated areas. Retreatment of this product may be necessary to obtain season long control.

MARGINAL WEEDS (Infesting the Edges and Other Portions of Ponds, Lakes, and Ditches):	GALS./100 GALS. WATER
Cattails (<i>Typha</i> spp.)	10

APPLICATION FOR MARGINAL WEEDS: For top kill, apply this product in 100 gallons of water plus a non-ionic surfactant* for full coverage and thorough weed control. Repeat treatment as necessary to control re-growth. For best results apply before flowering. *Add a Nonionic Surfactant containing 75% or greater surface active agent at the rate of 1/2 to 4 pints per 100 gallons of finished product.

APPLICATION FOR ALGAE: This product will control certain filamentous algae: Spirogyra spp. and Pithophora spp: Apply 0.5-1.5 ppm with product for control. Application methods as for submerged weeds.

Formula for Parts Per Million (ppm) Recommendations: Gallons to use - ppm recommended x acreage of water x average depth of water x 14. (Example: 1/2 ppm recommendations; two acre pond; 4 feet average depth; 1/2 x 2 x 4 x 14 = 56 gallons required.)

NOTICE: Buyer and user acknowledge and assume all risks and liability of use, storage, and handling of this product not in strict accordance with directions given herewith.

01/03

EPA REG. NO. 10807-202
EPA EST. NO. 10807-TX-1

MADE IN U.S.A. BY:

amreg
990 Industrial Park Drive
Marietta, GA 30062

Boo589

2 OF 2



2/2