

2/8/1988

212471 655
17

FEB - 8 1988

Ms. E. David Lewis
Pazianos Associates
1338 G Street, S.E.
Washington, D.C. 20003

Dear Mr. Lewis:

Subject: Terrazan PCNB Technical 96%
EPA Registration No. 10820-1
Terrazan 24%
EPA Registration No. 10820-4
Your Submission Dated January 15, 1988

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy of the label is enclosed for your records.

In your next printing the label must be revised to include the new statement "Do not contaminate water when disposing of equipment washwaters."

Sincerely yours,

Lois A. Rossi
Product Manager (21)
Fungicide- Herbicide Branch
Registration Division (TS-767C)

CONCURRENCES

SYMBOL							
SURNAME							
DATE							

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with labeling.

Do not plant root crops in FNB treated fields within 12 months of broadcast and banding applications unless FNB is registered for use on those crops.

Do not apply this product through any type of irrigation system.

BEANS (Root and Stem Rot - *Rhizoctonia solani*): **FURROW** - Application rates are to 14,500 linear feet of row for bush type beans and to 8,430 linear feet of row for pole beans. Apply 2-3 qts. in 8 to 10 gallons of water. Spray planting furrow and covering soil at planting time. Avoid applying directly to seed; other wise delayed emergence may occur. In California and Arizona apply 3-4 pints in 8 to 10 gals. of water. Apply in-furrow and incorporate with covering soil at planting. Use lower rate for lighter soils. Avoid application to bare seed.

SNAP and DRY BEANS (White Mold - *Sclerotinia sclerotiorum*): Use 1 gal. mixed in 8 to 10 gals. of water and apply to 14,500 linear feet of row for bush type beans or to 8,430 linear feet of row for pole beans. Spray in an 8" wide band centered on the row, immediately after or at time of seeding. Avoid applying directly to seed other wise delayed emergence may occur. (Where dinitro type pre emergence herbicides are used, add 1 gallon to the final herbicide spray mix and apply simultaneously. If disease is severe, application may be repeated at 2 to 3 week intervals using 1 or 2 nozzles per row and directing spray at base of plants. Do not apply after first bloom. Soil should remain undisturbed after treatment. Do not feed treated vines to livestock.

COTTON (Damping Off - *Rhizoctonia solani*): Use 1/2 to 1 gallon per acre in at least 15 gallons of water (per 13,000 linear feet of row based on 40" spacings). Spray on seed and surrounding soil at planting time. Do not allow the feeding or grazing of cotton foliage by livestock.


GARLIC (White Rot - *Sclerotium cepivorum*): Use 10 gals. mixed in 100 gals. of water and apply to 21,800 linear feet of row at planting time. Attach a suitable rig to the machine planter, using 2 nozzles per planting furrow. Direct the front nozzle to spray the bottom of the open furrow and the "seed" as it is dropped. Direct the rear nozzle to spray the covering soil.

ORNAMENTALS

AZALEA AND CAMELLIA (Azalea Petal blight and Camellia flower blight): Apply as a drench under plants at a rate of 1 1/2 qts. in 10 gallons of water per 150 sq. ft. Begin application prior to opening buds and repeat every 3 to 4 weeks through the bloom period.


LAWNS (Brown Patch - *Rhizoctonia solani*): For Southern or warm season lawns such as St. Augustine, Bermuda and similar grasses: Make one application in the fall or spring at first indication of infection. Mix 1 qt. in sufficient water (10-15 gals). To cover 1,000 sq. ft. Apply by means of a sprinkler can, spray or any other convenient method of obtaining even coverage. Following application, the treated area should be lightly watered to insure movement through the grass to the soil level. If the treated area is subjected to unusually heavy rainfall, or flooded, or if the disease is severe or reappears, the area should be retreated 3-4 weeks later. Under certain growing conditions, a temporary discoloration of the grass occasionally may occur. This causes no harm and will disappear in a short time. Do not graze treated areas. Do not feed clippings to livestock.

For Dichondra: Treat as above, using 1 1/2 qts. mixed in 40 gals. of water per 1,000 sq. ft.



TERRAZAN® 24%

EMULSIFIABLE CONCENTRATE



EPA REG. No. 10820-4

EPA EST. No. 10820 MX 01

ACTIVE INGREDIENT:	
Pentachloronitrobenzene (PCNB)	24.00 %
INERT INGREDIENTS	76.00 %
TOTAL	100.00 %

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed or inhaled. Avoid breathing spray mist. Use with adequate ventilation. Avoid contact with eyes, skin or clothing. Avoid contact with skin by wearing the following protective clothing (respirator, gloves, long-sleeved shirt, long pants, socks and shoes) must be worn during mixing and loading procedures. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Wear gloves and respirator when loading solvent. Avoid contamination of food and feed products.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, call a physician immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. If on skin, wash with plenty of soap and water. Get medical attention if irritation persists. If inhaled, remove victim to fresh air. Apply artificial respiration, preferably mouth to mouth, if indicated. If in eyes, flush with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not apply directly to water or wetlands (swamps, bays, marshes, and potholes). Do not contaminate water or wetlands by cleaning of equipment or disposal of wastes.

CHEMICAL HAZARDS

Flammable. Do not use or store near heat or open flame. Keep container closed.

STORAGE AND DISPOSAL


STORAGE: Store in a cool dry place. Protect from temperatures below 30° F.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.



MADE IN MEXICO BY



División Plásticos

Química Orgánica de México, S. A.

Am 115 Carretera a San Luis P. C. 2 P 21600 Mexicali B. C.
Adm Postal 3-1205 2 P 21100 Tels 6 8301 6106
P. O. Box 1840 Calexico Calif. 92231
Telex 056 9 881

282