

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed or inhaled. Avoid breathing spray mist. Use with adequate ventilation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Wear gloves and respirator when loading sprayer. Avoid contamination of food and feed products.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, call a physician immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
If on skin, wash with plenty of soap and water. Get medical attention if irritation persists.
If inhaled, remove victim to fresh air. Apply artificial respiration, preferably mouth to mouth, if indicated.
If in eyes, flush with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on this label.

CHEMICAL HAZARDS

Flammable. Do not use or store near heat or open flame. Keep container closed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store in a cool dry place. Protect from temperatures below 20°F.
PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.
CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.



Made in México by


Cydsa

ACTIVE
Per

INERT

TRIGRANTM-S

SEED PROTECTOR



ACCEPTED
with COMMENTS
In EPA Letter Dated:

DEC 6 1982

Under the Federal Insecticide, Fungicide, and Rodenticide Act, registration of this pesticide is required under EPA Reg. No. 10820-7

ACTIVE INGREDIENT:
 Pentachloronitrobenzene 24.7%

INERT INGREDIENTS 75.3%

Total 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. No. 10820-
EPA. EST. No. 10820 MIX 01

División Plásticos

Química Orgánica de México, S. A.
Km. 11.5 Carretera a San Luis P. C., Mexicali, B. C.
Apdo. Postal 3-1205 Tel. 8-79-01 al CE
P. O. Box 1840 Calexico, Calif. 92231

It is a violation in any manner inconsistent with the provisions of the Federal Seed Act.

GENERAL DIRECTIONS
Use solely for seed treatment to control fungus carried on seed and in the soils, desiccation, as well as reduction of seed losses.

Do not use for other agricultural purposes. Seed should be placed in contact with all applicable Seed Act. No other requirements.

APPLICATION

It is necessary to apply directly to the seed one to two equal parts of the table below

SEED	DISEASE
WHEAT	Cover
OATS	Cover
BARLEY	Cover
COTTON	Root Damp
BEAN (Snap & Dry)	Root White
CORN	Root
SOYBEAN	Root
SAFFLOWER	Damp Root

Mexico by

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL DIRECTIONS: TRIGRANTM has been developed solely for seed treatment to prevent the damage caused by fungus carried on seeds, or by fungus which already exists in the soils, destroying and damaging seed and seedlings as well as reducing the stands.

Do not use treated seed for food, feed or oil purposes. Seed treated with Trigran which is placed in containers must be labeled in accord with all applicable requirements of the Federal Seed Act. Note: some states have similar requirements.

APPLICATION RATES: Uniform application of TRIGRAN S is necessary to obtain adequate seed coverage and to produce the best results. TRIGRAN S may be applied directly to the seed (i.e., by drip method) or mixed in one to two equal parts of water and sprayed. Refer to the table below for appropriate seed treatment rates.

SEED	DISEASE	APPLICATION DOSAGE
WHEAT	Covered smut	2 oz./Bushel
OATS	Covered smut	2-4 oz./Bushel
BARLEY	Covered smut	2-4 oz./Bushel
COTTON	Root & Stem Rot Damping off	12 oz./100 lb (acid delinted) 16 oz./100 lb (delinted mechanically)
BEAN (Snap & Dry)	Root & Stem Rot White Mold	2 oz./Bushel
CORN	Root & Stem Rot	2 oz./Bushel
SOYBEAN	Root Rot	2-4 oz./Bushel
SAFFLOWER	Damping off Root Rot	2-4 oz./Bushel

NOTICE

Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals and condition of application equipment will vary, seller makes no claims other than those stated on this label.