

READ ENTIRE LABEL BEFORE USING

# DANGER POISON

DANGER

10825  
2

Toxaphene may be fatal if swallowed. Do not breathe spray mist. Do not get in eyes, on skin or clothing. If the concentrate spray is spilled on any part of the body or the clothing, wash thoroughly immediately. Use approved respirators, goggles and protective clothing. Keep away from children, domestic animals, feed or foodstuffs.

## PRECAUTIONS IN USING


Do not apply to fruits and vegetables after the edible portions begin to form or to forage crops which will be fed to dairy animals or to animals being finished for slaughter. Toxaphene is toxic to livestock and poultry and is very toxic to fish. This product is toxic to bees and should not be applied when bees are actively visiting the area.

This product is toxic to fish and wildlife. Keep out of lakes, streams, and ponds. Apply this product only as specified on this label. Do not apply when weather conditions favor drift of spray from areas treated. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies.

Birds feeding on treated areas may be killed.

## NON WARRANTY

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty



# TOP DOG BRAND

## 6 Lb. TOXAPHENE

### EMULSIFIABLE INSECTICIDE CONCENTRATE

## WARNING

Keep out of reach of children. Do not use in or near buildings or other structures. Do not use in or near water.

1 gallon TOP DOG 6 Lb. Toxaphene contains the equivalent of 6 Lbs. Toxaphene

## GENERAL I

COTTON: Dilute the amount of amount of water. Then add to water. To control boll weevils, cotton leafworm, fleahoppers, and concentrate per acre diluted with your equipment. For aerial and approximately 3 to 10 gallons per acre. On seedling cotton, 1/3 g usually sufficient with one nozzle. the plants become larger, increase concentrate per acre on grown cotton each row. Some burn may result therefore, it is advisable to have plants. For grasshoppers and crickets per acre. CAUTION - Cotton when this insecticide should not be being finished for slaughter.

## DIRECTION

OATS, RYE, WHEAT, BARN grasshoppers, chinch bugs and crickets use 1/3 gallon of the concentrate equipment application use appropriate dilute spray solution per acre. within 2 weeks of harvest. Do not weeks of harvest. Do not feed animals being finished for slaughter.

SOYBEANS For armyworm grasshoppers use 1/3 gallon of the ground equipment application use the total dilute spray solution per harvest. Do not feed treated plants to poultry, dairy animals slaughter. Do not feed soybean not apply more than 2/3 gallon application

ACCEPTED  
 6/27/72  
 LIN...  
 FOR...  
 FOR...  
 TO ATTAIN...  
 COMMENTS



**TOP  
 DOG  
 BRAND**

**6 Lb. TOXAPHENE**

**EMULSIFIABLE INSECTICIDE CONCENTRATE**

**WARNING**

KEEP OUT OF REACH OF CHILDREN  
 READ AND FOLLOW DIRECTIONS CAREFULLY

1 gallon TOP DOG 6 Lb. Toxaphene contains the equivalent of 6 Lbs. Toxaphene

**GENERAL DIRECTIONS**

**COTTON:** Dilute the amount of concentrate to be used with the same amount of water. Then add to this emulsion the required volume of water. To control boll weevils, thrips, bollworm, ~~fat~~ armyworm, cotton leafworm, fleahoppers, and grasshoppers, use 1/3 gallon of the concentrate per acre diluted with the necessary amount of water to fit your equipment. For aerial and ground equipment application, use approximately 3 to 10 gallons of the total dilute spray solution per acre. On seedling cotton, 1/3 gallon of the concentrate per acre is usually sufficient with one nozzle per row using ground equipment. As the plants become larger, increase the dosage to 1/2 gallon of the concentrate per acre on grown cotton, using not less than 3 nozzles to each row. Some burn may result on tender portions of the plant, therefore, it is advisable to have the nozzles at least 10 inches from the plants. For grasshoppers and crickets, use 1/3 gallon of the concentrate per acre. **CAUTION** - Cotton which has received late applications of this insecticide should not be grazed by dairy animals and animals being finished for slaughter.

**DIRECTIONS FOR USE**

**OATS, RYE, WHEAT, BARLEY, RICE:** For armyworms, grasshoppers, chinch bugs and climbing and surface feeding cutworms, use 1/3 gallon of the concentrate per acre. For aerial and ground equipment application use approximately 3 to 10 gallons of the total dilute spray solution per acre. Do not apply to oats, rye and wheat within 2 weeks of harvest. Do not apply to barley and rice within 3 weeks of harvest. Do not feed treated forage to dairy animals or animals being finished for slaughter.

**SOYBEANS:** For armyworms, leafhoppers, flea beetles and grasshoppers use 1/3 gallon of the concentrate per acre. For aerial and ground equipment application use approximately 3 to 10 gallons of the total dilute spray solution per acre. Do not apply within 21 days of harvest. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals, or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry. Do not apply more than 2/3 gallon of the concentrate per acre per application.

**ACTIVE INGREDIENTS (by weight)**

Toxaphene (Technical Chlorinated Camphene Chlorine content 67% to 69%)* .....	60%
<b>INERT INGREDIENTS:</b> .....	<b>40%</b>
<b>TOTAL</b> .....	<b>100%</b>

\* Technical Chlorinated Camphene (Chlorine content  
67% to 69%)

**DO NOT USE, POUR, SPILL OR STORE NEAR  
FLAME.**