

READ ENTIRE LABEL BEFORE USING

# DANGER POISON

## ANTIDOTE AND FIRST AID TREATMENT

**ATROPINE IS THE EMERGENCY ANTIDOTE FOR METHYL PARATHION POISONING.** Consult your physician about obtaining an adequate supply of 1/100-grain (0.6 mg.) atropine tablets for emergency use. 2-PAM is also antidotal and may be administered in conjunction with atropine. **CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.**

**INTERNAL:** If swallowed call a physician immediately. Induce vomiting by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of warm water). Repeat until vomit fluid is clear.

**NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.**

**EXTERNAL:** If the material has been spilled on the skin, immediately remove patient from the vicinity of the Methyl Parathion, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

**IF WARNING SYMPTOMS APPEAR:** Take or administer two 1/100-grain atropine tablets immediately. Get medical attention at once. 2-PAM is also antidotal and may be administered in conjunction with atropine. **NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR (See WARNING SYMPTOMS below).** Have a patient lie down and keep quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

## NOTE TO PHYSICIAN

**WARNING SYMPTOMS:** Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

**TREATMENT:** Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams, until complete atropinization has occurred. 2-PAM is also antidotal and may be administered in conjunction with atropine. Morphine is contraindicated. Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until the time for cholinesterase regeneration has been allowed as determined by blood test.

## DANGER

**POISONOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN. REPEATED INHALATION OR SKIN CONTACT MAY, WITHOUT SYMPTOMS, PROGRESSIVELY INCREASE SUSCEPTIBILITY TO METHYL PARATHION POISONING.**

Do not swallow or get in eyes, on skin, or on clothing. Do not breathe dust or vapors. Wear clean natural rubber gloves; clean, waterproof protective clothing, and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or

ACCEPTED  
7/21/72  
UNDER THE...  
FUNGIC...  
FOR LE...  
TO UNDER...  
TO ATTACHED COMMENT...



# TOP DOG BRAND

# 4 Lb. METHYL PARATHION

## EMULSIFIABLE INSECTICIDE CONCENTRATE

**DANGER POISON** KEEP OUT OF REACH OF CHILDREN  
SEE ANTIDOTE AND OTHER WARNINGS ON SIDE PANEL

1 gallon of TOP DOG BRAND 4 LB. METHYL PARATHION contains 4 pounds of Methyl Parathion

## PRECAUTIONS

Do not apply or allow to drift onto beneficial animals. Vacated areas of insecticide and volatile residues should be hand picked. Workers entering treated areas wear protective clothing. Do not slaughter in treated fields.

Highly toxic to fish and wildlife. Do not contaminate any body of water.

Do not apply where runoff is likely under conditions favor drift from areas. Do not contaminate food or feed. Surfaces to be decontaminated and destroyed should be removed to a safe place in a manner that contact with humans is avoided.

## GENERAL INFORMATION

**TOP DOG 4-lb. Methyl Parathion** is a low-volume ground applicator, recommended for use on all crops. Mix with water by agitation; agitate stand, re-agitate before use. Suggest

## DIRECTIONS

**INSECT CONTROL ON COTTAGE** control Aphids, Boll Weevil, Cotton Spider Mites.

4-lb. Methyl Parathion spray is recommended per acre depending upon local conditions. Consult your local representative of State Extension Service Specialists and Do not hand pick or harvest within

## DISPOSAL OF CONTAINER


**DO NOT** re-use the container. Do not use unused chemical at least 18 inches from supplies.

**FIVE GALLON AND FIFTY-FIVE GALLON DRUMS** - 2 qts. household detergent, 55-GALLON DRUMS - 1 cup detergent

Rotate container carefully until rinse solution at least 18 inches from supplies.

**DO NOT** use container in any case. (CAUTION: Handle caustic soda on skin, in eyes, or on clothing.) Punch holes in the top and bottom





# 4 Lb. METHYL PARATHION

## EMULSIFIABLE INSECTICIDE CONCENTRATE

### ACTIVE INGREDIENTS (by weight)

*O,O-dimethyl O-p-nitrophenyl thiophosphate	45.4%
Aromatic Petroleum Solvent .....	49.0%
INERT INGREDIENTS (by weight) .....	5.6%
TOTAL.....	100.0%

\*Methyl Parathion

WARRANTY AND LIMITATION OF DAMAGES. Willight Chemical Corp. warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein. Willight Chemical Corp. makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Willight Chemical Corp. is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or value, etc.

NOT FOR USE OR S  
DO NOT USE, POUR, SPI  
DO NOT USE THIS C