

PM 15
10827-56. 12/7/90

PRECAUTIONARY STATEMENTS

CAUTION HAZARDS TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED. MAY BE ABSORBED THROUGH THE SKIN. DO NOT BREATHE SPRAY MIST. AVOID CONTACT WITH SKIN, EYES, AND CLOTHING. WASH THOROUGHLY AFTER HANDLING. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. DO NOT USE ON HOUSEHOLD PETS OR HUMANS. DO NOT PERMIT CHILDREN OR PETS TO GO ON THE SPRAYED GRASS UNTIL THE SPRAY HAS COMPLETELY DRIED. KEEP OUT OF REACH OF DOMESTIC ANIMALS.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and wildlife. Keep out of lakes, streams, and ponds. Do not apply when weather favors drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are usually visiting to crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

Birds feeding on treated areas may be killed.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. PESTICIDE DISPOSAL: Pesticides, spray mixture or rinsate that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: 5, 20 & 35 gallon drums: a) Recycle container and offer for reconditioning, or b) triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved in 181, or bury in a safe place.

GENERAL: Consult Federal, State or local disposal authorities for approved alternative procedures.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

HOW TO MIX AND APPLY

NUMBER ONE

Determine size of spraying job. For lawns measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear, or in advance of an expected infestation. Repeat applications as needed or as noted in the rate chart.

NUMBER TWO

Use clean spray. Carefully measure and mix the amount of Diazinon and water as suggested in the chart.

NUMBER THREE

Spray as directed in the rate chart. Use coarse drizzling sprays for lawns, soil and other surface applications. Use medium or fine mist sprays for trees and plants. Thorough coverage is important. Flush sprayer with clean water after each use.

Nuisance Pests in Outside Areas

Ants, Boxelder Bug, Brown Dog Tick, Chigger, Clover Mite, Cockroach, Cricket, Earwig, Flea, Flea, Millipede, Spider, Springtails (Collembola).
16 fluid ounces in 3 gallons of water.

Thoroughly spray window and door screens, eave foundations, porch and patio walls and ceilings, entrance, walk, garbage cans, tree trunks, and into cracks and other places where insects can hide.

Do not treat plants or animals.
* Spray a 5 foot band of soil against the base next to the foundation of the house.

For Industrial & Commercial Use Only

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS

Formula 1250 EMULSIFIABLE CONCENTRATE

ACTIVE INGREDIENTS:

0,0-Diethyl 0-(2-isopropyl-6-methyl-4 pyrimidinyl) phosphorothioate*-----12.5%

INERT INGREDIENTS:

Aromatic Petroleum Solvent-----87.5%
TOTAL-----100.0%

Contains .96 lb. Diazinon per gallon

*DIAZINON® is a registered trademark of CIBA-GEIGY CORPORATION

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED — Call a physician immediately. Gastric lavage is indicated if material was taken internally. DO NOT INDUCE VOMITING unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED — Remove victim to fresh air. Apply respiration if indicated.

IF ON SKIN — Wash contaminated skin with soap and water.

IF IN EYES — Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

NOTE TO PHYSICIAN — If symptoms of cholinesterase inhibition are present, Atropine sulfate is indicated and may be administered with 2-PAM.

See side panel for additional precautionary statements.

Fire Ants

To aid in the control of fire ants in lawns and other recreation areas, apply 2 fluid ounces (4 tbsp.) in a minimum of 1 gal. of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal. per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness.

Note: For best results apply in cool weather, 66-80° F, or in early morning or late afternoon.

DIRECTIONS FOR USE (continued)

FOREIGN AREA

Pests — Rate and Area Covered

Lawns
Bent, Bermuda,
Bluegrass,
Biscuitgrass,
Fescue, Marion,
St. Augustine

Ants, Armyworms, Bermudagrass Weevil, Brown Dog Tick*, Chigger, Crickets, Cutworms, Earwigs, Fleas, Fruit Flies, Lawn Billbugs**, Chinch Bugs, Lawn Moths (Ear Webworms), Leafhoppers, Mites (Bermuda Clover), Sawflies (Billbugs), Springtails (Collembola).

2 fluid ounces in 3 gallons of water covers 125 square feet.

16 fluid ounces in 24 gallons of water covers 1,000 square feet.

Thoroughly wet down grass a few hours before applying.

* Brown Dog Tick: Do not apply to animals, in particular, spray under shrubbery and the grass next to houses.

** Lawn Billbugs: Water the lawn thoroughly immediately after treatment.

Note: In the North, if Chinch Bugs are the only lawn insect problem, a lower rate of application can be used.

1 fluid ounce in 3 gallons of water covers 125 square feet.

Millipedes: 4 fluid ounces in 3 gallons of water covers 125 square feet.

Thoroughly wet grass a few hours before applying.

Stinkgrass Fleas: 2 fluid ounces in 3 gallons of water covers 100 sq. ft. Apply when Sable Crawlers first emerge.

Ants, Biggus Weevil: 2 fluid ounces in 3 gallons of water covers 125 sq. ft. Spot spray ant hills and wasp ground nest openings.

Make the application at dawn when the weevils are not active.

Billbugs*, White Grubs** of Japanese Beetle, European Chaffer, Southern Chaffer.

2 fluid ounces in 3 gallons of water covers 125 square feet.

Water grass thoroughly after application.

* Billbug control: Apply when activity is first observed or when showed or brown grass is present, indicating damage from the insect.

** Grub control: Make application any time between late July and early October.

Hyperodes Weevil (New York only)

1 fluid ounce in 3 gallons of water covers 125 square feet.

Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

BEST AVAILABLE COPY

Trees & Shrubs
Arbutus,
Azalea, Birch,
Beverwood, Camellia,
Douglas Fir, Elm,
Hawthorn, Holly,
Juniper, Lilac,
Laurus, Maple,
Mimosa, Oak, Pine,
Plum, Poplar,
Rhododendron,
Spruce, Willow

Wet leaves and branches to the dripping point. Try to penetrate dense foliage.

Aphids, Bagworms, European Pine Shoot Moths, Holly Bud Moths, Dipterous Leaf Miners (Arbutus, Azalea, Birch, Beverwood, Holly, Oak), Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Two-spotted Spider), Oblique Root Weevils, Ominivorous Leaf Tiers, Scale Crawlers (Carnegiea, Eucalyptus, Laurus, Pine Needle, Ph. Scirp, San Jose, Saff),

2 fluid ounces in 3 gallons of water. Apply as a thorough foliar spray.

Webworms (Juniper, Mimosa), Whiteflies.

2 fluid ounces in 3 gallons of water. Apply as a thorough slow spray.

* Mimosa Webworms: Apply as a thorough foliar spray when adults first appear. Make two additional applications at 4-6 week intervals.

Apple-and-Thorn Skeletonizers, Humble, Charms, Oak Looper, Oblivious-banded leaf Rollers, Pear Slugs, Tom Caterpillars, Webworms, (Coronaster, Fall).

6 fluid ounces in 3 gallons of water. Apply as a thorough foliar spray.

Roses, Flowers
Azalea, Camellia,
Chrysanthemum,
Carnations,
Gladiolus, Lilac,
Rhododendron,
Roses

Wet the plants to a dripping point. Try to hit underside of leaves and penetrate dense foliage.

Aphids, Eucalyptus Scale Crawlers, Flea Beetles, Dipterous Leaf Miners, Leafhoppers, Mealybugs, Mites (Carnegiea Bud, Carnation Shear, Cyclamen, European Red, Two-spotted Spider), Thrips, Whiteflies.

2 fluid ounces in 3 gallons of water. Apply as a thorough foliar spray.

Do not use on certain forms, lilies, poinsettias, papaya, pine, geranium, and ornamentals since injury may occur.