

EVANS WOOD PRESERVATIVE

Oil base preservative that cleans up with water.

EVANS
PRODUCTS COMPANY
Roanoke, VA
24015

ACCEPTED
MAY 14 1982
U.S. Dept. of Agriculture
Forest Service
Forest Pest Control
Roanoke, VA 24015
EPA Reg. No. 10856-10

PHOTO OF
SIDING + DECK

NEW!
Oil base
preservative
that cleans up
with water.
Dries clear.


- Water repellent • Reduces swelling and shrinking
- Protects against mildew and rot • Dries clear.

ONE U.S. GALLON
3.785 LITERS

KEEP OUT OF REACH OF CHILDREN
WARNING
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

EPA Reg. No. 10856-10 EPA Est. No. 10856-VA-1
Active Ingredient
3-toda-2-propoxy-butyl carbamate 0.5%
Inert Ingredients 1.5%
Total 2.0%

WOOD PRESERVATIVE



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

- **HARMFUL IF SWALLOWED.** May be absorbed through skin. Causes skin irritation. Avoid breathing vapor or spray mist. Do not get in eyes, on skin, or on clothing.
- **FIRST AID STATEMENT.** If swallowed, do not induce vomiting. Call a physician immediately. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.
- **NOTE.** Wear goggles, rubber gloves and protective clothing when applying or handling treated lumber.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not add directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame. Use with adequate ventilation. Avoid breathing vapors. Never keep container inside home.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

STORAGE AND DISPOSAL

Do not reuse empty container. Wrap container and put in trash collect-on.

KEEP OUT OF REACH OF CHILDREN

EVANS CLEAR WOOD PRESERVATIVE—TYPE 48004

This non-flammable, timber pretreatment emulsion is designed to protect the wood surfaces around your home against mildew, rot and decay organisms. Any surface treated with this product may be easily painted as soon as it is dry.

Where to Use:

Recommended for use on millwork, plywood, decks, outdoor furniture, structural lumber, fences, shingles, siding—all exterior wood that may be subject to fungal attack. It is an ideal first coat prior to top-coating with paint or varnish.

Application:

For proper penetration, wood must be clean. Apply by brush, roller, spray or dipping. If immersion for three to five minutes is impractical, a second coat may be applied within one hour of the first coat. May be applied to green or damp wood. Not to be applied at temperatures of less than 40 F.

Coverage:

Approximately 100-300 square feet per gallon depending upon method of application and porosity of wood being treated.

Clean-Up:

Clean brushes or equipment with soapy water.

NOTE TO USER: Do not use on wood which will come into contact with food. Do not use on wood which will come into continuous contact with skin, unless the wood is sealed with paint, varnish or shellac after treatment. Do not use on wood in contact with the ground.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

May cause irritation of eyes, nose, throat and skin. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. In case of contact, flush with plenty of water. For eyes, get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL AND CHEMICAL HAZARDS

Keep away from fire and sparks. Store in a cool dry place.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Prohibitions and Recommendations: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

Keep away from fires and sparks. Store in a cool place, but not below 32° F. Do not allow to become overheated in storage. This may bring on chemical changes which will impair the fungicidal effectiveness of MF-4. Keep container closed when not in use.

Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

DIRECTIONS FOR USE AS A SPRAY

MF-4 is approved for application in sufficient water to provide good coverage with available equipment in either dilute sprays or in concentrate ground or aerial sprays. Rates listed are based on 100 gallons of dilute spray unless otherwise noted. Rates for concentrate and aerial applications should be comparable to the rates for dilute sprays on a per acre basis.

MF-4 may be applied by aircraft at the rate shown below for control of certain crop disease. See product labeling for listing of specific crops, details of timing, use limitations and other use instructions.

CROPS	Amount Per Acre	
	MF-4 (Qts.)	WATER (Gals.)
Tree Fruits & Nuts Vineyards	4 to 5.6	10 to 15
Potatoes and most vegetables;		
Small Fruits; Field Crops	1.2 to 1.6	5 to 8
Tomatoes, Sweet Corn	1.2 to 2.4	5 to 8

Apply with properly calibrated aerial equipment, arrange nozzles so that spray delivery is uniform over the entire spray swath.

AGR

ALMONDS: Brown Rot and petal fall or every 7 to 10 day intervals finished for slaughter. Do not apply later than 14 days before harvest.

APPLES: Black Rot, Sooty Blotch - 1.2 to 1.6 quarts at 7 to 14 day intervals; Arkansas, Delaware, Minnesota, Pennsylvania, South Carolina.

APRICOTS: Brown Rot full bloom, petal fall, and 1.6 quarts at full bloom. Do not apply later than 14 days before harvest.

BANANAS: Sigatoka - 1.6 quarts beginning at first signs of disease.

CRANBERRIES: Fruit rot - 1.6 quarts per 10 to 20, August 1 to 15 (Wisconsin) - 1.6 quarts within 30 days of harvest.

FIGS (Kadota): Sooty Blotch 20 days before harvest.

GRAPES: Black Rot again 20 days later; do not apply later than 14 days before harvest.

NECTARINES: Brown Rot, and at 7 to 14 day intervals.

PAPAYAS: Anthracnose (not less than 50.0 gallons fruit is set. Repeat at 10 day intervals to central column).

PEACHES: Brown Rot dormant period. Do not apply later than 14 days before harvest.

AG

ASPARAGUS: Rust Repeat at 10 day intervals bears and repeat at 10 day intervals.

BEANS (Lima): Anthracnose Do not apply later than 14 days before harvest.

BEANS (Green and Broad): Repeat at 5 to 7 day intervals.

BROCCOLI: Alternaria Mildew begin in plant before harvest. Remove residue.