



# TIFCHEM

## WETTABLE SULPHUR

### GUARANTEED ANALYSIS

ACTIVE INGREDIENTS: Sulphur, not less than 90%  
INERT INGREDIENTS: Not more than 10%

(Sulfur fineness 93% through 325 mesh screen.)

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH INSTRUCTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS: WE RECOMMEND REFERENCE TO AGRICULTURAL EXPERIMENT STATIONS ON SPECIFIC USAGE.

**Caution:** Keep out of reach of children

### DIRECTIONS

CITRUS: Rust Mite -- Use 5 to 10 lbs. per 100 gallons water. During cool weather use 10 lbs. of TIFCHEM Wettable Sulphur per 100 gallons water.

PEACHES: Brown Rot, Powdery Mildew, Scab -- 4 to 6 lbs. per 100 gallons water. Make applications during pink, petal fall and cover spray periods. Also as ripening of fruit approaches for Brown Rot Control.

APPLES: Apple Scab, Powdery Mildew -- 6 to 8 lbs. per 100 gallons water. Make application between periods of rainfall. Repeat as necessary to cover new foliage growth during pink, petal fall and cover spray periods.

CAUTION: Some fruits and other plants are susceptible to injury from sulphur under certain climatic conditions. The user is advised to not use sulphur on any crop unless local use has proven that sulphur is safe in that locality. During periods of high temperature sulphur may burn foliage and fruit. Do not make sulphur applications at excessively high temperatures. Do not use TIFCHEM Wettable Sulphur with oil or within 3 weeks after oil application.

### NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.

**DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.**

TIFTON CHEMICAL COMPANY  
Tifton, Georgia

EPA Reg. No. 10873-3

**CAUTION:** Keep Out Of Reach Of Children. Harmful if swallowed. Do not breathe spray mist. Wash thoroughly after using. Do not re-use container. Destroy when empty.

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.** Buyer assumes all risks of use, storage or handling. This material not in strict accordance with label directions.

**SHADE TREES, FLOWERS, SHRUBS;** (Azaleas, Begonias, Camellias, Chrysanthemums, Crepe Myrtle, Dahlias, Delphinium, Fuchsias, Roses, Snapdragons, Stocks, Sweet Peas) - Black Spot of Rose, Leaf Spots, Powdery Mildew, Rust: Mix 1 Tbsp. per gallon water. Spray when foliage or disease appear. Repeat at 7-10 day intervals or as necessary.

**APPLES, PEARS - Powdery Mildew:** Mix 1 Tbsp. per gallon water. Spray when buds swell. Repeat when fruit buds show color and again at petal fall.

**APRICOTS - Shothole:** Mix 1 Tsp. per gallon of water. Apply in delayed dormant period (during January in West Coast areas). Repeat at 7-10 day intervals as necessary.

# *Hacienda*

## **LIQUID**

# **Copper Fungicide**

## **SPRAY**

### **ACTIVE INGREDIENTS**

Copper Oleate*	15.0%
Xylene	79.4%
<b>INERT INGREDIENTS</b>	<b>5.6%</b>

\*Copper expressed as elemental 1.34%

## **NET CONTENTS 1 GALLON**

NECTAR  
Mildew  
(Nov. 1)  
to swell  
WALNUT  
after bl  
vals as  
CITRUS  
Spray 1  
1 Tbsp.  
as nec  
BLACK  
Powder  
disease  
TOMATO  
Spray b  
necessa  
BEANS  
Spray b  
necessa  
CUCUM  
Begin v  
necessa  
MELON  
melon,  
Tbsp. p  
7-10 da