


1081  
P-21 S  
441-4

# TIFCHEM

## 3.4-70 Peanut Dust



[FUNGICIDE]

NET WEIGHT 50 LBS.

Active Ingredients	By Wt.
Copper (in mixture of cuprous and cupric oxides) expressed as metallic	3.4%
Sulfur	70.0%
Inert Ingredients	26.6%
Sulfur Fineness Through 325 Mesh—93% Typical	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

This product is formulated for application by ground or air equipment.

### DIRECTIONS

**PEANUTS:** Leaf Spot Infections—Apply 2 to 25 pounds of TIFCHEM 3.4-70 Dust per acre at 7 to 10-day intervals or as necessary. Begin dusting when diseases first appear and continue to within 1 month of harvest.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

**CAUTION:** Harmful if swallowed. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use have proved that sulfur does not damage crops in that locality. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

ACCEPTED

NOV 12 1971

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
ED UNDER NO. 16873-5

**CONDITIONS OF SALE:** 1. **TIFTON** Chemical Company (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond the manufacturer's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above) resulting from handling, storage and use of this material.

TIFTON CHEMICAL COMPANY  
Tifton, Georgia

EPA REG. NO. 10873-5

Active Ingredients		By Wt.
Copper (in mixture of cuprous and cupric oxides) expressed as metallic .....		3.4%
Sulfur .....		70.0%
Inert Ingredients .....		26.6%
Sulfur Fineness Through 325 Mesh—93% Typical.		

**READ ENTIRE LABEL. USE STRICTLY IN ACCORD-  
ANCE WITH CAUTIONS, WARNINGS AND DI-  
RECTIONS, AND WITH APPLICABLE STATE AND  
FEDERAL REGULATIONS.**

**KEEP PESTICIDE IN ORIGINAL CONTAINER.**

**BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.**

**This product is formulated for application by  
ground or air equipment.**

### **DIRECTIONS**

**PEANUTS: Leaf Spot Infections**—Apply 20 to 25  
pounds of **TIFCHEM 3.4-70** Dust per acre at 7 to 10-  
day intervals or as necessary. Begin dusting when  
diseases first appear and continue to within 1 month  
of harvest.

**CAUTION:** Harmful if s  
and other plants are suscepti  
fur under certain climatic co  
advised not to use sulfur on  
use have proved that sulfur d  
in that locality. Do not apply  
volving possible drift to fo  
plantings that might be do  
thereof rendered unfit for sal

**TIFTON CHEMICAL**  
**Tifton, Geo**