

ACCEPTED

NOV 17 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND FUNGICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 10873-6
SUBJECT
To Attached COMMENTS

10873-6

NET WEIGHT 50 LBS.

Active Ingredients	By Wt.
Carbaryl (1-naphthyl N-methylcarbamate)	10.0%
Copper (In mixture of cuprous and cupric oxides) expressed as metallic	3.4%
Sulfur	70.0%
Inert Ingredients	16.6%

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS; WE RECOMMEND REFERENCE TO AGRICULTURAL EXPERIMENT STATIONS ON SPECIFIC USAGE.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS

PEANUTS: Blister Beetles, Mexican Bean Beetle, Early and Late Leaf Spot Infections—10 to 15 lbs. per acre. Thrips, Leafhoppers, Three-cornered Alfalfa Hopper, Cucumber Beetles, Bean Leaf Beetle, Japanese Beetle, Alfalfa Caterpillar, Velvet Bean Caterpillar, Armyworm, Webworms,

TIFCHEM

SEVIN* Copper-Sulfur 10-3.4-70 PEANUT DUST

CAUTION

READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid Prolonged or Repeated Breathing of Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Contact. Wash Hands and Face Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily. Note for Physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal. AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

*Trade Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate

BEST DOCUMENT AVAILABLE

TIFTON CHEMICAL COMPANY

Tifton, Georgia

Corn Earworm, Stink Bugs, Grasshoppers—15 lbs. per acre. May be applied up to the day of harvest or grazing. Begin application at first sign of insects or disease and repeat at regular intervals as necessary to maintain control.

NOTICE ON CONDITIONS OF SALE: 1. TIFTON Chemical Company (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty, guarantee or representation, express or implied, concerning this material.

2. Because critical, unforeseeable factors beyond the manufacturer's control prevent it from eliminating all risks in connection with the use of chemicals even though reasonably fit for such use, buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above) resulting from handling, storage, and use of this material. These risks include, but are not limited to, damage to plants, crops and animals to which the material is applied, failure to control pests, damage caused by drift to other plants or crops, and personal injury. Buyer and user accept and use this material on these conditions, whether or not such use is in accordance with directions.

EPA REG. NO. 10873-6

7

NET WEIGHT 50 LBS.

Active Ingredients	By Wt.
Carbaryl (1-naphthyl N-methylcarbamate)	10.0%
Copper (In mixture of cuprous and cupric oxides) expressed as metallic	3.4%
Sulfur	70.0%
Inert Ingredients	16.6%

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS; WE RECOMMEND REFERENCE TO AGRICULTURAL EXPERIMENT STATIONS ON SPECIFIC USAGE.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS

PEANUTS: Blister Beetles, Mexican Bean Beetle, Early and Late Leaf Spot Infections--10 to 15 lbs. per acre. Thrips, Leafhoppers, Three-cornered Alfalfa Hopper, Cucumber Beetles, Bean Leaf Beetle, Japanese Beetle, Alfalfa Caterpillar, Velvet Bean Caterpillar, Armyworm, Webworms,

KEEP OUT OF REACH OF CHILDREN

CAUTION: HARMFUL IF INHALED OR ABSORBED THROUGH SKIN. Avoid Prolonged or Repeated Breathing of Vapors. Take Internally. **Skin Contact:** Avoid Contact. Wash Hands and Face Thoroughly. Take Shower or Bath After Work. Change and Wash Sleeved Work Clothing. Change Daily. **Note for Physician:** Carbaryl is a reversible, Cholinesterase inhibitor. Atropine is an antidote. AVOID CONTAMINATION OF FEED, FEEDING TROUGHS AND WATER TACLES. Some fruits and other plants are susceptible to injury from sulfur under certain conditions. The user is advised not to use on any crop unless local use has been shown to do no damage crops in that area. Sulfur is highly toxic to bees. For information on treatment or residues on crops, contact your local Extension Station. For more information may be obtained from the Agricultural Extension Service. Use on conditions involving possible damage to other plantings that might be rendered unfit for consumption.

TIFTON CHEMICAL COMPANY

Tifton, Georgia