

UPPER PANEL FOR INSTRUCTIONS
 LOWER PANEL FOR PRECAUTIONS
 SEE SIDE PANEL FOR OTHER WARNINGS

WARNING

- MAY BE FATAL IF SWALLOWED.
- DO NOT BREATHE SPRAY MIST.
- MAY BE ABSORBED THROUGH SKIN.
- DO NOT GET IN EYES, ON SKIN, OR ON CLOTHING.
- AVOID CONTAMINATION OF FEED AND FOOD-STUFFS
- WASH THOROUGHLY AFTER USING.

DO NOT reuse container.
 Destroy when empty.

NOTICE

Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk to person or property arising out of use or handling and accepts the product on these conditions.

WARNING

This product is toxic to bees and should not be applied when bees are actively visiting the area.

TIFCHEM

TOXAPHENE EM-6

Emulsifiable Liquid
 (Contains 6 lbs. Toxaphene Per Gal.)
AN AGRICULTURAL INSECTICIDE

ACTIVE INGREDIENTS:	
Toxaphene*	58.0%
Xylene-Range Aromatic Hydrocarbon Solvent	37.0%
INERT INGREDIENTS:	5.0%
TOTAL	100.0%
*Technical Chlorinated Camphene (Chlorine Content 67% to 69%)	

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME!
 PROTECT FROM TEMPERATURES BELOW 0° F.

WARNING: KEEP OUT OF REACH OF CHILDREN.
 SEE SIDE PANEL FOR OTHER WARNINGS

TIFTON CHEMICAL COMPANY
 Tifton, Georgia

Pour recommended
 Keep agitator running
 mixture does not
 result. DO NOT
 previous use of the

**OBSERVE
 USE**

DO NOT use this p
 fed to dairy anim
 for slaughter.

Highly toxic to fis
 when weather co
 treated. Do not
 occur. Do not co
 equipment or disp

DO NOT use on c
 melon, squash) a

Observe use limita
 specific crops.

The Federal Food a
 official tolerances
 In order that res
 will not exceed
 if recommended
 than specified.

Consult the State A
 additional inform
 and rate of app
 directions for vary

CHEM

HENE EM-6

Emulsifiable Liquid
(5.0% Toxaphene Per Gal.)
AGRICULTURAL INSECTICIDE

.....	58.0%
Hydrocarbon Solvent	37.0%
.....	5.0%
.....	100.0%
Toxaphene (Chlorine Content 67% to 69%)	

KEEP AWAY FROM NEAR HEAT OR OPEN FLAME!
TEMPERATURES BELOW 0° F.

KEEP OUT OF REACH OF CHILDREN.

SEE LABEL FOR OTHER WARNINGS

AGRI-CHEMICAL COMPANY

Waynesboro, Georgia

DIRECTIONS FOR USE

Pour recommended amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing, and spraying operations. DO NOT allow mixture to stand. If mixture does not mix readily, but tends to separate as an oily layer, DO NOT use as injury to plants may result. DO NOT combine emulsifiable liquids with wettable powders in the same spray tank unless previous use of the materials being combined has proved them to be physically compatible.

OBSERVE THE FOLLOWING

USE PRECAUTIONS

DO NOT use this product on feed or forage to be fed to dairy animals or to animals being finished for slaughter.

Highly toxic to fish and wildlife. DO NOT apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

DO NOT use on cucurbits (cantaloupe, cucumber, melon, squash) as injury to plants may result.

Observe use limitations given on the label for specific crops.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates and DO NOT apply higher than specified.

Consult the State Agricultural Extension Service for additional information on the timing, number, and rate of applications consistent with label directions for varying local conditions.

RECOMMENDATIONS

Dosages are given in terms of pints per acre. Use in sufficient water for thorough coverage (Approximately **100-125 gallons**).

For airplane application, use approximately **3 to 10 gallons** of water per acre, consistent with crop growth and good coverage.

COTTON: For control of Armyworms, Bollworms, Cotton Bollworms, Grasshoppers, apply 1-1/3 quarts per acre. DO NOT graze dairy animals or animals being finished for slaughter on cotton or cotton trash.

EPA Reg. No. 10873-10

_____ **GAL. NET**