


TIFCHEM

2% PHOSDRIN® DUST



* If you are not a U.S. citizen, please indicate your country of origin: _____

[illegible]

POISON



MINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR

WARNING SYMPTOMS. Symptoms include weakness, headache, tachycardia, blurred vision, dry mouth, distended pupils, salivation, sweating, nausea, vomiting, diarrhea, and abnormal cramps. **TREATMENT:** Atropine is the specific therapeutic antagonist of, hence, against parasympathetic stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be given to rectify hemodynamic effects and doses of 1 to 2 mg/kg, intravenously, until complete antiparasympathetic effect is obtained. ECG should be obtained and may be administered for 24 hours with atropine. Monitor for contrast material leak, chest hypoxia, myocardial infarction, and other complications. Observe patient and record vital signs, repeat exposure, and pulse rate and blood pressure with atropine. If prolonged symptoms occur, give small doses of any of the above antidotes. All who further experience symptoms should be rechecked and may be allowed as frequently as desired.

DIRECTIONS

APPLICATION A large number of applications have been submitted for consideration by the Commission. The following are some of the more recent ones:

SORGHUM (*Sorghum vulgare*)—A grain crop native to Africa, sorghum has been grown in the United States since the early 1800s. It is well adapted to hot climates and drought conditions. Sorghum is used for animal feed, human consumption, and industrial purposes.

BEANS

BEETS BROCCOLI BRUSSELS SPROUTS CABBAGE CAULIFLOWER COLLARDS, KALE, LETTUCE, MUSTARD, SPINACH, TURNIPS

[illegible]

GRAPES *Vitis* spp. are woody vines that are cultivated for their fruit. Although many varieties are grown for wine, the most common are the seedless table grapes. The fruit is eaten fresh or dried into raisins. Grapes are grown in many regions of the world, but the United States is the largest producer and consumer. The fruit is a cluster of small berries. The skin is usually green or purple, and the flesh is white or green. The seedless varieties are called 'seedless' or 'parthenocarpic'. They are produced by a process called 'parthenocarpy', which is the development of fruit without the need for pollination. This is achieved by the use of plant hormones or by selective breeding. The seedless grapes are often marketed as 'Jumbo' or 'Super' sizes. They are popular for their sweet taste and ease of eating. The seedless varieties are also used for raisins. The seedless grapes are often grown in California, where they are the most common variety. They are also grown in other parts of the world, including Europe, China, and India. The seedless grapes are a popular choice for many people, especially children, because they are easy to eat and do not contain seeds. They are also a good source of vitamins and minerals. The seedless grapes are often sold in bunches, and they are a popular choice for gifts. They are also a popular choice for parties and events. The seedless grapes are a delicious and healthy fruit that is enjoyed by many people around the world.

At the end of the study, the authors concluded that the results of the study suggest that the use of the proposed model can be used to predict the effect of the use of the proposed model on the performance of the system. The authors also concluded that the use of the proposed model can be used to predict the effect of the use of the proposed model on the performance of the system.

**DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING
WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES**

TIFTON CHEMICAL COMPANY
ORREST CITY, ARKANSAS -- TIFTON, GEORGIA

EPA Reg. No. 10873-14

TIFCHEM

2% PHOSDRIN® DUST

Aug. 25-1972

44

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 10573-150 POUNDS

10F1
Pm 16
44 7-13

DANGER! KEEP OUT OF REACH OF CHILDREN

 **POISON** 

ATROPINE IS THE EMERGENCY ANTIDOTE FOR PHOSPHORUS INSECTICIDE POISONING. If you suspect that you or someone else has been poisoned with phosphorus insecticide, **CALL A PHYSICIAN IMMEDIATELY** in all cases of suspected poisoning. **INTERNAL** symptoms may include nausea, vomiting, diarrhea, and abdominal pain. **EXTERNAL** symptoms may include redness, swelling, and pain at the site of contact. **NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.** If the person has been exposed to the insecticide, remove contaminated clothing and wash the exposed area with soap and water. **IF BURNING SYMPTOMS APPEAR**, wash the area with large amounts of water. **NEVER ADMINISTER ATROPINE UNLESS BURNING SYMPTOMS APPEAR.** Symptoms may be relieved by giving the person 1 to 2 mg of atropine. If symptoms persist, repeat the dose. **NEVER ADMINISTER ATROPINE TO A CHILD WITHOUT THE ADVICE OF A PHYSICIAN.**

WARNING SYMPTOMS: Symptoms include weakness, restlessness, tightness in chest, blurred vision, nonreactive or pinpoint pupils, tachycardia, sweating, nausea, vomiting, diarrhea, and abdominal cramps. **TREATMENT:** Atropine in the special therapeutic antagonism of this drug against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10-minute intervals in doses of 1 to 2 mgm orally, until complete atropinization has occurred. If ECG is also abnormal, and may be administered in conjunction with atropine. Atropine is also contraindicated in fear chest by postural damage. Oxygen administration may be necessary, thus the patient continuously, for 10 hours. Repeated exposure to this contraindication is not allowed, without warning. The following procedure should be followed: Give small, slow, every 30 minutes as much as 10 mgm. No further treatment until time for the next dose. If necessary, a physician should be called for further treatment.

PRECAUTIONS IN USING: This product is toxic to fish and wildlife. Birds and other wildlife untreated areas may be killed. Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor runoff from treated areas. Treat only nonpotable water by cleaning off equipment, and disposal of wastes. Apply this product only as specified on the label. Keep out of the reach of children and domestic animals. Do not consume raw food from this product. Employees are not to be destroyed by burning or burning. It is flammable and explosive. Do not use in hand equipment, and do not use in the presence of open flames. Do not use in the presence of heat. Apply only when the label first appears to appear and do not use in the presence of heat as necessary.

Begonia application: When insects first appear and repeat as often as necessary to maintain control.

GRAPES: Apply at the rate of 12 to 25 pounds per acre for control of Aphids, and 25 to 50 pounds to control Leaf Folder, Leafhoppers, Mites, Red-banded Leafhopper, and Lygus Bug. Do not apply within 2 days of harvest.

STRAWBERRY: Apply at the rate of 6 to 12 pounds per acre for control of Aphids and Mites, and 12 to 25 pounds per acre for control of Grasshoppers, Strawberry Leafhopper, Salt-marsh Caterpillar, and Lygus Bug. Do not apply within 1 day of harvest.

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.

SORGHUM Apply to sorghum in permanent culture at 10 to 25 pounds for control of corn rootworm and corn root borer. Apply for control of corn rootworm and corn root borer in sorghum at 10 to 25 pounds.

[illegible]

NOTICE: The directions for use given for this product are based upon use under normal and reasonably foreseeable conditions. If the directions for use are fol-

circumstances, if, however, because of the nature of the commodity, geographical location, and other conditions under which this product may be used, it is clearly evident to the seller that there are no warranties of representation, express or implied, concerning this material and its use except that it conforms to the chemical description on the label and no person is authorized to make any representation on upon its behalf, the seller shall not be held responsible for any damages resulting from the handling, storage or use of this product for any reason in connection with its use of the material, conform to the chemical description on the label, damages, if any, which may result from the failure of this material to conform with the description on the label shall be limited to direct damages but shall in no case consequential

EPA REG. NO. 16873-14