

CRICKET BAIT

Contains 5% **DYLOX**

ACTIVE INGREDIENT:

Dimethyl (2,2,2-trichloro-1-hydroxyethyl) Phosphonate 5.00%

INERT INGREDIENTS 95.00%

Total 100.00%

DYLOX is a Reg. TM of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen.

KEEP OUT OF REACH OF CHILDREN

CAUTION

May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use. Do not contaminate feed or food. If misuse results in poisoning, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression. TO PHYSICIAN: Atropine sulfate is antidotal. 2-Pam is also antidotal and may be administered in conjunction with atropine.

Manufactured By:

TIFTON CHEMICAL COMPANY

P. O. Box 5, Tifton, Georgia 31794

NET CONTENTS 50 LBS.

EPA Est. No. 10873-GA-1

EPA REG. NO. 10873- 29

ACCEPTED
NOV 14 1975
 for the Federal Insecticide, Fungicide, and Rodenticide Act, amended, for the pesticide registered under EPA Reg. No. 10873-19

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Crop	Insect	Rate/Acre	Details of application and comments
Corn	Army worm Cricket Saltmarsh caterpillar Surface feeding cutworms	20 lb.	Apply as a band over the row, around the base of plants or as a broadcast treatment. Apply before or just after planting fields with a history of surface feeding cutworm damage and to those fields suspected of being infested. Do not apply within 40 days of harvest. Do not apply more than once per season.
Cowpeas (black-eyed peas, crowder peas, southern peas)	Army worm Cricket Saltmarsh caterpillar Surface feeding cutworms	30 lb.	Apply as a band over the row, around the base of plants or as a broadcast treatment. Apply before or just after planting fields with a history of surface feeding cutworm damage and to those fields suspected of being infested. Do not apply within 14 days of harvest. Do not apply more than twice per season. Do not feed treated vines to dairy or meat animals.
Tobacco	Army worm Cricket Saltmarsh caterpillar Surface feeding cutworms	20 lb.	Apply as a band over the row, around the base of plants or as a broadcast treatment. Apply before or just after planting fields with a history of surface feeding cutworm damage and to those fields suspected of being infested. Do not apply after plants are 2 feet tall.
Peanuts	Army worm Cricket Saltmarsh caterpillar Surface feeding cutworms	20 lb.	Apply as a band over the row, around the base of plants or as a broadcast treatment. Apply before or just after planting fields with a history of surface feeding cutworm damage and to those fields suspected of being infested. Make up to 3 applications before digging. One additional treatment may be made between digging and harvest.

or
 the
 int
 with
 im-
 ing
 of
 bod