

20%

TIFCHEM  
TOXAPHENE

DUST

## DIRECTIONS FOR USE

To be used as a dust without dilution in conventional ground and air dusters. Adequate and uniform coverage of the plant is essential. Dusts should be applied when the wind velocity is less than 5 M.P.H. The best hours for dusting are usually between 5:00 P.M. and 9:00 A.M. Apply only to foliage when needed unless otherwise noted in special instructions.

CROP & INSECTS	RATE PER ACRE	INTERVAL Application to Harvest
COTTON***—Fleashoppers, Thrips, Grasshoppers	5-10 lbs.	
Boll Weevil, Bollworm, Cotton Leafworm, Cutworms (Climbing & Surface), Fall Armyworm, Lygus and Other Mirids, Rapid Plant Bug, Yellow-striped Armyworm, Flea beetles	10-25 lbs	
Garden Webworm	20-30 lbs	
Do not apply after the bolls open. Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.		

BEANS—Armyworms, Serpentine Leaf Miner, Corn Earworm, Cutworms (Climbing & Surface), Thrips, Lygus Bugs, Compsa, Curculio, Salt-marsh Caterpillar, Leafhoppers, Grasshoppers, Daring Ground Beetle, Bean Lycaenid

10-25 lbs. 7 days

No limitation on use of shelled beans as human food. Do not feed treated vines to dairy animals or animals being finished for slaughter.

BROCCOLI, BRUSSEL SPROUTS, CABBAGE, CAULIFLOWER—Armyworms, Cabbage Looper, Cutworms (Climbing & Surface), Diamond Back Moth Larvae, Imported Cabbage Worms, Corn Earworm, Thrips

10-35 lbs.

Do not apply after edible parts begin to form or within 30 days of harvest

CARROTS—Cutworms (Climbing & Surface), Grasshoppers

10-20 lbs. No Limitations

CELERY—Armyworms, Cutworms (Climbing & Surface), Thrips, Serpentine Leafminer

10-20 lbs.

Do not apply after plants start to bunch or after plants are half mature

COLLARDS, KALE—Cutworms (Climbing & Surface), Thrips, Serpentine Leaf Miner, Armyworms, Cabbage Worm, Green Peach, Aphid

10 lbs. 14 days

Cabbage Looper

10-20 lbs. 21 days

EGGPLANT, PEPPERS, PIMENTO\*\*—Colorado Potato Beetle, Flea beetle, Potato Aphid, Leafhopper, Armyworms, Serpentine Leafminer, Blister Beetle

10-15 lbs. 5 days

\*\*10 lbs. \*\*2 days

HORSE RADISH, PARSNIPS, RUTABAGAS—Striped Flea beetle, Grasshoppers, Flea beetles

7½-15 lbs.

Do not use tops for food or feed

LETTUCE—Cutworms (Climbing & Surface), Armyworms

10-15 lbs.

Do not apply after seedling stage on leaf lettuce, mustard greens, Swiss Chard, or after heads start to form on head lettuce. Do not apply within 7 days of harvest on head lettuce if outer leaves are to be stripped. Do not apply after heads begin to firm if outer leaves are not to be stripped

ONIONS—Thrips

10-15 lbs.

Do not apply to green or spring onions.

NET CONTENTS 50 LBS.

## ACTIVE INGREDIENTS:

\*Toxaphene ..... 20%

INERT INGREDIENTS ..... 80%

Total ..... 100%

Technical Chlorinated Camphene (Chlorine content 67 to 69%)

KEEP OUT OF REACH OF CHILDREN

## CAUTION

Harmful if swallowed. Avoid Breathing Dust. Avoid contact with eyes, skin and clothing. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Do not treat areas where livestock feed or graze.

Avoid applications that will carry residue on edible portions of crops when harvested for market. Do not use on dairy animals or dairy barns.

## ENVIRONMENTAL HAZARDS

"This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where there are important resources. Apply this product only as specified on this label."

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning (Keep out of smoke).

This product is toxic to bees and should not be applied when bees are actively visiting area.

SPINACH—Cutworms (Climbing & Surface), Thrips 10 lbs 21 days  
Serpentine Leaf Miner, Armyworms, Cabbage Worm, Green Peach Aphid, Cabbage Looper  
Do not apply more than once per season

TIFTON CHEMICAL CO.

TIFTON, GEORGIA

TOMATOES—Tomato Fruitworm, Serpentine Leafminer, Pinworm, Tomato Hornworm, 10 lbs. 1 day  
Tobacco Hornworms, Blister Beetles, Southern Green Stink Bug, Cutworms (Climbing & Surface), Armyworms, Stink Bugs 15-25 lbs. 3 days  
Tomato Russet Mite 25 lbs. 3 days  
CRANBERRIES—Armyworms Grasshoppers 10-25 lbs.

Do not apply after the fruit starts to form.

STRAWBERRIES—Strawberry Crown Borer, Strawberry Weevil, Cutworms (Climbing & Surface)

10-15 lbs.

Do not apply after the fruit starts to form.

BARLEY, RYE, WHEAT—Cutworms (Climbing & Surface), Armyworms Grasshoppers 10-15 lbs.

Do not feed treated forage to dairy animals or animals being finished for slaughter. No restriction on the use of the grain.

CORN—Fall Armyworm, Corn Earworm, Grasshoppers, Budworm, Cutworms (Climbing & Surface), Lantern Fly, Chinch Bug, Alfalfa Webworm 10-30 lbs.

Do not feed treated forage to dairy animals or animals being finished for slaughter. No restriction on the use of the grain.

PEANUTS—Cutworms (Climbing & Surface) Fall Armyworms, Velvet Bean Caterpillar, Budworm 10-30 lbs.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

SORGHUM—Armyworms, Budworm, Corn Lantern Fly, Chinch Bugs, 10 lbs. 28 days  
10-15 lbs. 40 days  
Do not apply more than once after heads start to form. Do not ensile treated forage. Do not graze dairy animals or animals being finished for slaughter on treated fields.

SOYBEANS—Velvet Bean Caterpillar, Bean Leaf Beetle, Grasshoppers, Corn Earworm, 10-20 lbs. 21 days  
Fall Armyworms, Thrips, Blister Beetle  
No limitations on the use of threshed beans. Do not apply less than 3 weeks before harvest. Do not feed treated plants or ensilage made from treated plants to poultry or dairy animals, or animals being finished for slaughter.

TOBACCO (Plant Beds only)—Tobacco 7½-10 lbs.  
Cutworms  
Apply 7-10 days before transplanting.

## NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.

EPA REG. NO. 10873-21  
EST. NO. 10873-GA-1