

TIFCHEM SEVIN* 5 Dust

[INSECTICIDE]

NET WEIGHT

Active Ingredient	By Wt.
Carbaryl (1-naphthyl N-methylcarbamate)	5%
Inert Ingredients	95%

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

This product is formulated for application by ground or air equipment.

ALFALFA, CLOVERS, COWPEAS, PEANUTS, SOYBEANS: Blister Beetles, Mexican Bean Beetle—10 to 20 lbs. per acre. Alfalfa Caterpillar, Bean Leaf Beetle, Cucumber Beetles, Green Cloverworm, Japanese Beetle, Leafhoppers, Three-cornered Alfalfa Hopper, Thrips, Velvetbean Caterpillar—20 lbs. per acre. Armyworms, Corn Earworm, Stink Bugs, Webworms—20 to 30 lbs. per acre. May be applied up to harvest. Some leaf injury may occur if applications are made when tender foliage is wet.

APPLES: Codling Moth, Orange Tortix (Skin Worm)—Use 80 lbs. per acre. Do not use within one day of harvest.

APRICOTS: Codling Moth, Peach Twig Borer, Leafroller, — 75 to 80 lbs. per acre. Do not apply within 3 days of harvest.

BEANS: Mexican Bean Beetle—10 lbs. per acre. Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Velvet Bean Caterpillar, Flea Beetles, Japanese Beetle—20 lbs. per acre. Corn Earworm, Armyworms, Tarnished Plant Bug, Stink Bugs—20 to 30 lbs. per acre. In California for Corn Earworm, Lima Bean Pod Borer, Lygus and Stink Bugs—40 lbs. per acre. May be applied up to harvest.

BLUEBERRIES: Blueberry Maggot, Cherry and Cranberry Fruitworms, Japanese Beetles—30 to 40 lbs. per acre. Apply 3 weeks before harvest and repeat 10 days later or as necessary. May be applied up to harvest.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI: Flea Beetles, Harlequin Bug—10 to 20 lbs. per acre. Armyworms, Corn Earworm, Imported Cabbageworm, Young Cabbage Looper, Salt-Marsh Caterpillar—20 to 40 lbs. per acre. Do not apply within 3 days of harvest.

CORN: Corn Earworm. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations. Corn Rootworm Adults, European Corn Borer, Fall Armyworm, Sap Beetles, Japanese Beetle, Leafhoppers, Flea Beetles—30 to 40 lbs. per acre. For insects attacking silk and ears apply at 2 to 5 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. May be applied up to harvest.

CRANBERRIES: Cutworms, Fireworms, Fruitworms, Japanese Beetle, Leafhoppers—30 to 40 lbs. per acre. Apply in late bloom and as needed at 7 to 10-day intervals. Do not apply within 1 day of harvest.

CUCUMBERS, MELONS, PUMPKINS, SQUASH: Pickleworm, Melonworm—10 to 20 lbs. per acre. Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs—20 lbs. per acre. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use SEVIN on watermelons in Florida. May be used up to harvest.

Aug 31 1973
 UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN.

CAUTION: HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN, Avoid Breathing of Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Contact. Wash Hands and Face Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily. Note for Physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal. AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

OKRA: Corn Earworm, Stink Bugs—20 to 40 lbs. per acre. Repeat at 5 to 7-day intervals or as necessary.

PEACHES: Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Cat-Facing Insects, Periodical Cicada—70 to 80 lbs. per acre. Do not apply within 1 day of harvest.

POTATO, EGGPLANT, PEPPER: Colorado Potato Beetle, Flea Beetles, Leafhoppers—10 to 20 lbs. per acre. Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Lygus Bugs, Stink Bugs, Lace Bugs, Tarnished Plant Bugs—20 to 40 lbs. per acre.

RICE, FORAGE GRASSES, PASTURE: Armyworms, Stink Bugs, Thrips—20 to 30 lbs. per acre. For optimum thrip control in grasses grown for seed, direct dust into boot. Do not apply to Rice within 14 days of harvest. May be applied up to harvest on Pastures and Forage Grasses.

SORGHUMS (Including Milo and Grain Sorghum): Armyworms, Corn Earworm, Stink Bugs, Webworms—20 to 40 lbs. per acre. Sorghum Midge—30 lbs. per acre. Dust into the forming heads. Treatment for sorghum midge control should be made 3 to 4 days after heads have emerged from boot. Do not apply to Sorghum Grain within 21 days of harvest.

STRAWBERRIES: Meadow Spittlebug, Strawberry Leaf Roller, Strawberry Weevil—20 to 40 lbs. per acre. Do not apply within 1 day of harvest.

SUGAR BEETS: Armyworms, Leafhoppers, Webworms—20 to 40 lbs. per acre. Do not apply within 14 days of harvest.

TOMATO: Flea Beetles, Leafhoppers, Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer—50 to 60 lbs. per acre.

TOBACCO (In Plant Beds—except Kentucky): Tobacco Flea Beetle—Apply 1 to 2 lbs. per 100 sq. yds. when plants appear. Repeat at 10 to 14-day intervals and just before transplanting. (In Fields—All Areas)—Broadcast Applications—Hornworms, Tobacco Flea Beetles—Use 1 to 25 lbs. per acre depending on size of plants. Use higher dosage on large plants and repeat as needed. Good coverage is essential. Heavier applications may cause plant injury. If late applications are necessary, treat immediately after pruning. Do not apply within 5 days of priming or 15 days of cutting. Avoid getting wet with plant juices when cutting the crop.

POULTRY (Chickens, Partridges, Pheasants, Pigeons): Northern Fowl Mite, Lice and as a supplement to premise treatments for Chicken Mite, Fleas—1 lb. per 100 birds. Apply thoroughly by means of a shaker can or hand duster. Repeat in 4 weeks if necessary. Do not use on Poultry or game birds within 7 days of slaughter. **IN OR AROUND POULTRY HOUSES:** Chicken Mite, Fleas, Bed Bugs—1 lb. per 40 sq. ft. of litter or floor space, and to roosts and adjacent areas by means of a shaker can, hand duster or power duster. Force dust into cracks and crevices. Repeat as necessary. Do not treat nest litter. Avoid direct contamination of eggs. Do not use in premise treatments within 7 days of slaughter.

DOGS, CATS: Brown Dog Ticks, Fleas—Dust liberally over the animal and rub thoroughly into the skin. Apply also to the feet, between the toes and on the legs. Apply at weekly intervals if needed. Do not treat kittens or puppies under four weeks of age.

IN OR AROUND DOG AND CAT QUARTERS: Ticks, Fleas—Dust liberally in and around sleeping quarters, kennels and dog houses by means of a shaker can, hand or power duster. Force dust into cracks and crevices. Repeat as necessary but not more often than once weekly.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

*Trade Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate.

MANUFACTURED BY:

TIFTON CHEMICAL COMPANY

EPA REG. NO. 10873-24