

ENVIRONMENTAL HAZARDS

PRECAUTIONS IN USING: "This product is toxic to fish and wildlife," "Birds and other wildlife in treated areas may be killed," "Keep out of lakes, streams, or ponds," "Do not apply where runoff is likely to occur," "Do not apply when weather conditions favor drift from areas treated," "Do not contaminate water by cleaning of equipment, or disposal of wastes," "Apply this product only as specified on this label."

BEE CAUTION: This product is toxic to bees and should not be applied when bees are actively visiting the area.

RESTRICTIONS

USE PRECAUTIONS: Do not apply to or allow to drift onto varieties of sorghum which are sensitive to phosphates. To do so may cause burning of the sorghum or milo. Do not use on other crops used for food or forage. Use only according to label directions. Application at rates above those shown may result in illegal crop residues. Do not treat food crops grown in the greenhouse.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TIFCHEM CUTWORM & CRICKET BAIT

Contains 5% **DYLOX**

ACTIVE INGREDIENT:

Dimethyl (2,2,2-trichloro-1-hydroxyethyl) Phosphonate† 5.00%

INERT INGREDIENTS 95.00%

Total 100.00%

DYLOX is a Reg. TM of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen.

KEEP OUT OF REACH OF CHILDREN

CAUTION

May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot

Lima Bean
Cabbage
Cauliflower
Pepper
Collard
& Lett
Tomato
Crop
Corn
Cowpea
(black
crow
sour
Tobacco
Peas

BEE CAUTION: This product is toxic to bees and should not be applied when bees are actively visiting the area.

RESTRICTIONS

USE PRECAUTIONS: Do not apply to or allow to drift onto varieties of sorghum which are sensitive to phosphates. To do so may cause burning of the sorghum or milo. Do not use on other crops used for food or forage. Use only according to label directions. Application at rates above those shown may result in illegal crop residues. Do not treat food crops grown in the greenhouse.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

RECOMMENDATIONS

Crop	Insect	Rate/Acre	Details of application and comments
Beans (Dry & snap)	Cricket Saltmarsh caterpillar Westim bean cutworm Variegated cutworm Surface feeding cutworms (including black cutworm granulate cutworm	30 lb	Apply as a band over the row around the base of plants or as a broadcast treatment. Apply before or just after planting fields with a history of surface feeding cutworm damage and to those fields suspected of being infested. Do not apply within 14 days of harvest. Do not use treated vines as food for dairy or meat animals.

CRICKET BAIT

Contains 5% **DYLOX**

ACTIVE INGREDIENT:

Dimethyl (2,2,2-trichloro-1-hydroxyethyl) Phosphonate 5.00%

INERT INGREDIENTS 95.00%

Total 100.00%

DYLOX is a Reg. TM of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen.

KEEP OUT OF REACH OF CHILDREN

CAUTION

May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use. Do not contaminate feed or food. If misuse results in poisoning, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression. TO PHYSICIAN: Atropine sulfate is antidotal. 2-Pam is also antidotal and may be administered in conjunction with atropine.

Manufactured By:

TIFTON CHEMICAL COMPANY

P. O. Box 5, Tifton, Georgia 31794

NET CONTENTS 50 LBS.

EPA Est. No. 10873-GA-1

EPA REG. NO. 10873- 29

TIFCHEM CUTWORM & CRICKET BAIT

Contains 5% **DYLOX**

ACTIVE INGREDIENT:

Dimethyl (2,2,2-trichloro-1-hydroxyethyl)
Phosphonate† 5.00%

INERT INGREDIENTS 95.00%

Total 100.00%

DYLOX is a Reg. TM of the Parent Company of
Farbenfabriken Bayer GmbH, Leverkusen.

KEEP OUT OF REACH OF CHILDREN

CAUTION

May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use. Do not contaminate feed or food. If misuse results in poisoning, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.

Lima Beans	Amy worm Saltmarsh caterpillar Cricket Surface feeding cutworms Western bean cutworm	30 lb	Apply as a band over the row, around the base of plants or as a broadcast treatment. Apply before or just after planting fields with a history of surface feeding cutworm damage and to those fields suspected of being infested. Do not apply within 30 days of harvest. Do not apply more than twice per season.
Cabbage Cauliflower Peppers	Amy worm Saltmarsh caterpillar Cricket Surface feeding cutworms	20 lb	Apply as a band over the row, around the base of plants or as a broadcast treatment. Apply before or just after planting fields with a history of surface feeding cutworm damage and to those fields suspected of being infested. Do not apply within 21 days of harvest.
Collards & Lettuce Tomatoes	Amy worm Cricket Saltmarsh caterpillar Surface feeding cutworms	20 lb	Apply as a band over the row, around the base of plants or as a broadcast treatment. Apply before or just after planting fields with a history of surface feeding cutworm damage and to those fields suspected of being infested. Do not apply within 28 days of harvest.

Crop	Insect	Rate/Acre	Details of application and comments
Corn	Amy worm Cricket Saltmarsh caterpillar Surface feeding cutworms	20 lb	Apply as a band over the row, around the base of plants or as a broadcast treatment. Apply before or just after planting fields with a history of surface feeding cutworm damage and to those fields suspected of being infested. Do not apply within 40 days of harvest. Do not apply more than once per season.
Cowpeas (black-eyed peas, crowder peas, southern peas)	Amy worm Cricket Saltmarsh caterpillar Surface feeding cutworms	30 lb	Apply as a band over the row, around the base of plants or as a broadcast treatment. Apply before or just after planting fields with a history of surface feeding cutworm damage and to those fields suspected of being infested. Do not apply within 14 days of harvest. Do not apply more than twice per season. Do not feed treated vines to dairy or meat animals.
Tobacco	Amy worm Cricket Saltmarsh caterpillar Surface feeding cutworms	20 lb	Apply as a band over the row, around the base of plants or as a broadcast treatment. Apply before or just after planting fields with a history of surface feeding cutworm damage and to those fields suspected of being infested. Do not apply after plants are 2 feet tall.
Peanuts	Amy worm Cricket Saltmarsh caterpillar Surface feeding cutworms	20 lb	Apply as a band over the row, around the base of plants or as a broadcast treatment. Apply before or just after planting fields with a history of surface feeding cutworm damage and to those fields suspected of being infested. Make up to 3 applications before digging. One additional treatment may be made between digging and harvest.

ACCEPTED