



**TIFCHEM** PTED

OCT 22 1975  
Under the Federal  
Fungicide, Insecticide,  
and Plant Growth Regulator Act  
as amended, registered under  
EPA Reg. No. 10873-42

**USE NO HOOKS  
KEEP WHERE DRY**

# TOXAPHENE

## 40% WP

**AN INSECTICIDE FOR FORMULATING USE**

**ACTIVE INGREDIENT:**

Toxaphene (technical chlorinated camphene—  
67.69% chlorine)..... 40%

**INERT INGREDIENTS:**..... 60%

**WARNING:** May be fatal if swallowed. ● Do not get on skin, in eyes, or on clothing. ● Wash thoroughly after using. ● Can be absorbed through skin. ● Do not breathe dust.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. ● Do not contaminate water by cleaning of equipment or disposal of wastes. ● Rinse equipment and dispose of wastes by burying in noncrop lands away from water supplies. ● Do not reuse containers. Destroy when empty. ● Avoid contamination of feed and foodstuffs.

**ANTIDOTE:**

**INTERNAL:** Give sodium chloride 15 g (one tablespoonful of table salt) in a glass of warm water and repeat until vomit is clear. Call a doctor immediately.

**EXTERNAL:** Wash with soap and water.

**NOTE:** For additional information on the properties, handling, and use of Tifchem Toxaphene 40% WP refer to Tifchem Product Data Bulletin Number 102.

EPA Est. No. 10873—GA—1

**50.7 POUNDS • 23.0 KILOS GROSS WEIGHT**  
**50.0 POUNDS • 22.7 KILOS NET WEIGHT**


**PHYSICIAN'S NOTE:** Toxaphene produces central nervous system excitation that responds to barbituric acid derivatives. Prevent or control convulsions with barbiturates, preferably administered intravenously. If the barbiturate can be given before convulsions start, sodium pentobarbital is the drug of choice. Since toxaphene has a fairly long duration of action, a long-acting barbiturate such as phenobarbital may be used after initial control of convulsions. Refer to "Clinical Handbook of Economic Poisons," which was prepared by the U. S. Department of Health, Education, and Welfare, Public Health Service, Communicable Disease Center, Atlanta, Georgia.

**WARNING: KEEP OUT OF REACH OF CHILDREN**  
**NOTICE**

Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg. No. 10873—42

Manufactured By  
**TIFTON CHEMICAL COMPANY**  
Tifton, Georgia



# COASTOX META CAL 3-5 PELLETS

### ACTIVE INGREDIENTS:

Tricalcium Arsenate .....	5.00%
Metalddehyde .....	3.00%
INERT INGREDIENTS .....	92.00%
	<hr/>
	100.00%

Arsenic Expressed as metallic ..... 1.88%  
 Arsenic in water soluble form, expressed as metallic, not more than .50%



## POISON

## WARNING



# KEEP OUT OF REACH OF CHILDREN

THIS PESTICIDE MAY BE FATAL TO CHILDREN AND DOGS OR OTHER PETS IF EATEN.  
 KEEP CHILDREN AND PETS OUT OF TREATED AREA.

POISONOUS, HARMFUL IF SWALLOWED. AVOID BREATHING DUST. AVOID CONTACT WITH SKIN, EYES OR CLOTHING. AVOID STORAGE NEAR OR CONTAMINATION OF FEED, FOODSTUFFS OR DRINKING WATER. FIELD WORKERS SHOULD USE PROTECTIVE GLOVES WHEN HANDLING AND BROADCASTING META CAL SNAIL PELLETS. WASH WITH SOAP AND WATER AFTER HANDLING AND BEFORE EATING OR SMOKING.

IMPORTANT: Protect life and domestic animals. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

CALL A PHYSICIAN IMMEDIATELY. Until he arrives follow the directions here given. Contains ARSENIC (Tricalcium Arsenate)

ANTIDOTE: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Then two tablespoonfuls of Epsom Salt or Milk of Magnesia in water and force fluids. Have patient lie down and keep warm.

### USAGE CAUTIONS

Do not apply on food crops or directly to foliage or against the base of plants. Injury may result from bait becoming lodged in plant crevices. Birds and other wildlife feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not contaminate water by cleaning of equipment, or disposal of wastes and containers. Apply this product only as specified on the label.

Keep children and animals out of treated areas. Dogs may be attracted to this bait. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

### RECOMMENDATIONS

COASTOX META CAL 3-5 SNAIL PELLETS is a special pelletized formulation of special attractants including apple pomace plus active ingredients calcium arsenate and metalddehyde. After careful blending and mixing, the product is put through a pellet mill to give a uniform pelleted product. COASTOX META CAL 3-5 SNAIL PELLETS are recommended for the control of cutworms, slugs, and snails.

FOR SLUGS, SNAILS, CUTWORM CONTROL in ornamental gardens, nurseries in non-crop areas. Scatter pellets at the rate of 5 to 10 lbs. per 1000 sq. ft. Apply by hand or power driven equipment which is capable of directing the bait to the soil surface. Protective gloves should be worn when handling and broadcasting the bait. Do not contaminate the edible portion of any food crops. If soil is dry, sprinkle lightly. Best results are obtained in the evening.

FOR SLUG, SNAIL, CUTWORM CONTROL IN THE FOLLOWING BUSH AND VINE FRUITS (BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES, STRAWBERRIES, RASPBERRIES). Use 100 to 300 pounds per acre or 3 lbs. to 6<sup>3</sup>/<sub>4</sub> pounds per 1000 sq. ft. Apply to soil surface around the plant. Do not contaminate edible parts.

FOR SLUG, SNAIL, CUTWORM CONTROL IN AVOCADOS AND CITRUS (ORANGES, LEMONS, LIMES, GRAPEFRUIT, TANGELOS, TANGERINES). Use 50 pounds maximum per acre. Broadcast under trees. Do not apply to tree foliage on edible parts.

FOR SLUG, SNAIL, CUTWORM CONTROL IN VEGETABLE CROPS: ASPARAGUS, BEANS, BLACKEYED PEAS, BROCCOLI, BRUSSEL SPROUTS, CABBAGE, CANTALOUPE, CARROTS, CAULIFLOWER, CELERY, COLLARDS, CORN, COWPEAS, CUCUMBERS, EGGPLANTS, KALE, KOHLRABI, MELONS, PEPPERS, PUMPKINS, RUTABAGAS, SPINACH, SQUASH (summer), SQUASH (winter), TOMATOES, TURNIPS, WATERMELONS: Use 3 pounds to 6<sup>3</sup>/<sub>4</sub> pounds per 1000 sq. ft. or 100 to 300 pounds per acre. Apply to soil surface around the plants. May be applied by power driven or hand operated spreaders but do not allow pellets to contaminate the foliage on edible portions of any food or feed crops.

## CAUTION

Do not apply directly to foliage or against the base of plants. If bait becomes lodged in plant crevices burning may result. If bait is applied near a food crop, extreme care should be exercised so that there will not be any residue at harvest time.

NOTICE TO USER: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA EST NO 11656CA03 EPA NO 11656-22 # 40B