

**PRECAUTIONARY
STATEMENT**

Hazardous to Humans (and
Domestic Animals)

CAUTION

Reasonable care should be exer-
cised to avoid inhalation or con-
tact with open wounds.

**ENVIRONMENTAL
HAZARDS**

Keep out of lakes, streams or
ponds. Do not contaminate water
by cleaning or by disposal in
waters.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of federal law to use this
product in a manner inconsistent with its
labeling.

**STORAGE AND
DISPOSAL**

STORAGE:

Reclose containers of unused
DIPEL 150 Dust. Store in a dry
place.

DISPOSAL:

Do not reuse container. Discard
empty container according to
local trash disposal regulations.

NET CONTENTS 25 LBS.



DIPEL® 150 DUST

(Bacillus thuringiensis)

ACTIVE INGREDIENTS

Bacillus thuringiensis, Berliner, Potency of 320 Inter-
national Units per mg. (at least 0.5 billion viable spores
per g.)* 0.064%

INERT INGREDIENTS 99.936%

*Equivalent to potency of 0.15 billion International
Units per pound of this product.

• Dipel is the registered trademark of Abbott Laboratories,
North Chicago, Illinois

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel for Additional Precautionary Statements.

Manufactured By:

TIFTON CHEMICAL COMPANY

P.O. Box 5—Tifton, Georgia 31794

EPA Est. No. 10873-GA-1

EPA Reg. No. 10873-52

CROPS

DIPEL 150 Dust is a biological insecticide
intended for use against chewing larvae
(caterpillars) of certain lepidopterous
species.

DIPEL 150 Dust is recommended for use
by ground or by air. Apply when pests are
first noticed and repeat as necessary.
Thorough plant coverage is necessary for
control.

APPLICATION RATE

For the control of Cabbage Looper on
Broccoli, Cabbage, Cauliflower, Collards,
Celery, Lettuce, Melons, Potatoes,
Spinach, Tomatoes and Cucumbers
(Southeastern U.S. only) apply 25 to 50
lbs. per acre (approximately 1/2 - 1
lb./1000 sq. ft.), equivalent to 3.75 to 7.5
billion International units per acre.

For the control of Imported Cabbage
Worm on Broccoli, Cabbage, Cauliflower,
Collards, and Turnip Greens, apply 15 to
25 lbs. per acre (approximately 1/2 - 1/2
lb./1000 sq. ft.), equivalent to 2.25 to
3.75 billion International units per acre.

For the control of Hornworm on Tobacco
apply 12 to 25 lbs. per acre (approximate-
ly 1/4 - 1/2 lb./1000 sq. ft.), equivalent to
1.875 to 3.75 billion International units
per acre.

For the control of Grape leaf folder on
grapes, apply 25 to 50 lbs. per acre, (ap-
proximately 1/2 to 1 lb./1000 sq. ft.)
equivalent to 3.75 to 7.5 billion Interna-
tional units per acre.

WARRANTY STATEMENT

Seller makes no warranty, expressed or
implied, concerning the use of this pro-
duct other than indicated on the label.
Buyer assumes all risk of use and/or
handling of this material when such use
and/or handling is contrary to label in-
structions.