

ADCO MIST SPRAY INSECTICIDE

USDA Reg. No. 10885-1

Active Ingredients 100.00%
 *Piperonyl Butoxide Technical 0.75%
 Pyrethrins 0.15%
 Petroleum Distillate 99.10%

*Equivalent to .60% (butylcarbityl) (6-propylpiperonyl) ether and .15% related compounds.

CAUTION: Keep Out Of Reach Of Children

HARMFUL IF SWALLOWED. AVOID CONTAMINATION OF FOODS. DO NOT SPRAY ON HUMANS. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Use ADCO Mist Spray Insecticide as a space spray in food processing plants only when plant is not in operation. Foods and food products should be covered or removed during treatment. All food processing surfaces should be covered, or thoroughly cleaned, before resuming operations.

ADCO CHEMICALS, INC.

1307 Kirkland Dr.

Raleigh, N. C. 27603

Net Contents

..... Gals.

DIRECTIONS FOR USE

IN FOOD PLANTS, GRAIN MILLS, BARN AND ANIMAL SHELTERS

1. **FLIES, MOSQUITOES, AND GNATS:** Close windows and doors. Shut off ventilating fans. Direct spray toward upper portion of building. Fill entire room with mist. Keep area closed at least 10 to 15 minutes. Sweep up and destroy fallen insects. **Do not remain in treated areas and ventilate the area after treatment is completed.**
2. **ROACHES, WATERBUGS AND SILVERFISH:** Spray directly on insects, hitting as many as possible, while working the spray into all cracks, crevices, under and behind equipment, cabinets, sinks and other hiding areas. Thorough and repeated spraying is needed to help break insect breeding cycle.
3. **SPIDERS:** Direct wet spray to hit spiders and saturate spider webs and adjacent cracks and crevices.
4. **WEEVIL, FLOUR BEETLES, CONFUSED FLOUR BEETLES AND FLOUR MOTHS** in accessible or exposed stages. Spray walls, floors, partitions, bins, empty containers in flour mills, feed mills and processing plants with enough solution to wet the surfaces and penetrate all cracks and crevices and other insect hiding areas. Also, direct spray to hit all exposed ends, tops and sides of storage bags. Repeat treatment on a regular basis in warm weather insect migration periods.
5. **STABLE FLIES AND HORN FLIES:** Apply as a space spray mist with animals on premises. For temporary protection of dairy cattle and beef cattle from fly annoyance direct mist spray above backs of animals, using about one ounce and not more than two ounces per animal. Do not spray on udders of milk animals. Udders should be washed with a sanitizing solution before milking.
6. **FLEAS IN PREMISES:** Spray infested area liberally, working well into floor covering materials, floor cracks, baseboards and wall cracks and crevices. Thorough and repeated treatment of animal sleeping quarters is recommended. *XII indoors edible*

XII indoors non-edible **ACCEPTED**

MAY 3 1971

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. *10885-1*