


10900-63

4-29-2003

1110

Please read instructions on reverse before completing form.

Form approved OMB No. 2070-0060 Approval expires 05-31-98

EPA United States Environmental Protection Agency Washington, DC 20460		Registration Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number 245241
Application for Pesticide – Section I			
1. Company/Product Number 10900-63		2. EPA Product Manager Marion Johnson	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) 876 Institutional Insecticide		PM# 10	
5. Name and Address of Applicant (Include ZIP Code) Sherwin-Williams Diversified Brands 601 Canal Road Cleveland, OH 44113 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c) (3) (b) (I), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	
Section – II			
<input type="checkbox"/> Amendment – Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification – Explain below.		Final printed labels in response to Agency letter dated “Me Too” Application. Other – Explain below.	
NOTIFICATION APR 29 2003			
Explanation: Use additional page(s) if necessary. (For section I and Section II) Notice of label change relative to PR-Notice 2001-6. This notification is consistent with the guidance in PR Notice 2001-6 and the requirements of EPA's regulations at 40 CFR 156.10 and 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the guidance of PR Notice 2001-6 and the requirements of 40 CFR 156.10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.			
Section – III			
1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If “Yes” Unit Packaging Wgt. No. per Container	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If “Yes” Unit Packaging Wgt. No. per Container	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify)
*Certification must be submitted			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other			
Section – IV			
1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Jamie Zajac		Title Regulatory Affairs Specialist	Telephone No. (Include Area Code) (216) 515-8892
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punished by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature 		4. Title Regulatory Affairs Specialist	
4. Typed Name Jamie Zajac		5. Date 04/16/03	

2/10

876
Industrial
Insecticide

NOTIFICATION

APR 29 2003

ACTIVE INGREDIENTS:

*•+Resmethrin ([5-(phenylmethyl)-3-furanyl]
methyl 2,2-dimethyl-
3-(2-methyl-1-propenyl)
cyclopropanecarboxylate..... 0.2%w/w

**d-trans Allethrin..... 0.4%

OTHER INGREDIENTS:..... 99.4%
100.0%w/w

* Cis/trans isomers ratio:

max. 30%(+) cis and min. 70%(+) trans

• U.S. Patent Nos. 3,465,007 and 3,542,928.

**Bioallethrin

EPA Reg. No. 10900-63

EPA Est. No.

10900-OH-1
33915-MI-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for additional precautionary statements.

20 oz.(1 lb. 4 oz.) * 566 g net wt.

READ ENTIRE LABEL BEFORE EACH USE
USE ONLY ON DOGS OR CATS

A Synthetic pyrethroid for controlling the following flying & crawling insects in and around households, greenhouses, buildings, vehicles, boats, kennels & stables:

Flying Insects

houseflies
 mosquitoes
 gnats(midges)
 deer flies
 stable flies
 flying moths
 (millers)
 fruit flies
 wasps
 hornets
 yellow jackets
 bees

Crawling Insects

cockroaches
 waterbugs
 bedbugs
 fleas
 ticks
 ants
 crickets
 silverfish
 earwigs
 spiders
 sowbugs
 centipedes
 millipedes
 grain mites
 grain insects

Plant Pests

aphids
 thrips
 whiteflies
 leafhoppers
 two-spotted mites
 plant bugs
 army worms
 cutworms

leaf beetles
 gypsy moths
 tussock moths
 oakworms
 sawflies
 tent caterpillars
 red spider mites

spittle bugs
 weevils
 Japanese beetles
 slugs
 mimosa webworm
 spider mites
 budworms

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior landscapes or ornamental gardens.

For use indoors and outside premises of hotels, motels, homes, preschools, children's nurseries, day camps, noncommercial greenhouses, lathouses, and seed houses, commercial, industrial, office, institutional and storage buildings, milk handling areas of milk plants, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, flour and feed mills, bakeries, boats, ship cabins and hospitals. Also for use in campers, mobile homes, buses, automobile, trucks, boxcars and ship cargo holds. Also for use in dog kennels and horse stables. In horse stables, not for application if horses will be used for food.

4/10

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Shake well before use. Remove protective cap. Aim spray opening away from person. Push button to spray.

Food Areas: Food should be removed or covered during treatment. All food processing surfaces, shelving, utensils, etc., should be covered or removed prior to treatment or thoroughly washed with an effective cleaning compound followed by a potable water rinse prior to use. When using this product in these areas, apply only when the facility is not in operation.

Indoors: For use in household, commercial and institutional buildings, hospitals, noncommercial greenhouses and lathouses, campers, mobile home, vehicles and boats, as listed above. In hospitals, remove patients prior to treatment. Ventilate rooms for two hours before returning patients.

To Kill Flying Insects: houseflies, mosquitoes, gnats (midges), deer flies, stable flies, flying moths (millers), fruit flies. Close all doors and windows. Point nozzle upward. Direct spray mist to all parts of the room, especially window and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed. Repeat as necessary.

To Kill Wasps, Hornets, Yellow Jackets, Bees: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places, especially under attic rafters, contacting as many insects as possible. Treat these areas at the rate of two seconds per linear foot at a distance of 12 to 18 inches. Repeat as necessary.

To Kill Crawling Insects: Fleas, cockroaches, waterbugs, crickets, silverfish, earwigs, spiders, sowbugs, centipedes, millipedes, grain mites and grain insects (adult and accessible stages of confused flour beetles, red flour beetle, sawtoothed grain beetles, shiny spider beetles, carpet beetle, grain weevils, Angoumois grain moths, Indian meal moths). Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, storage areas and other insect harborage areas. Spray directly on insects where possible. Treat these areas at the rate of two seconds per linear foot at a distance of 12 to 18 inches. Repeat as necessary.

Directions for use continued:

To Kill Grain Insects: For control of the insect pests listed above, remove infested grain, cereal or flour and destroy. Avoid contamination of other food and foodstuffs in the area by removing before spraying: Spray directly on exposed stages and into cracks and crevices where grain insects may be hiding. In homes, dispose of all infected products. The shelves and drawers should be cleaned and thoroughly sprayed so as to contact all pests, and covered with shelf paper after allowing shelves to dry. Treat these areas at the rate of two seconds per linear foot at a distance of 12 to 18 inches. Any utensils that are contacted with spray should be washed before using. Repeat as necessary.

To Kill Spiders: Spray directly on these pests and use repeated applications to the webbing and all cracks and crevices which harbor spiders. Spray at the rate of two seconds per linear foot at a distance of 12 to 18 inches. Repeat as necessary.

To Kill Ants: Spray doors, around window frames, and trails and hills. Spray at the rate of two seconds per linear foot at a distance of 12 to 18 inches. Repeat as necessary.

To Kill Bedbugs: Apply as a spot treatment to cracks and crevices around baseboards, floorboards, bedboards and walls. First, thoroughly clean and air mattresses and springs. Then spot treat only areas that may harbor bedbugs, concentrating on tufts, folds and seams. Treat these areas at the rate of two seconds per linear foot at a distance of 12 to 18 inches. Repeat as necessary. For persistent infestations, consult a professional applicator.

To Kill Fleas & Ticks on Dogs & Cats: Apply directly to animals along back, under forelegs, and other areas except scrotum and anus. Part hair to contact skin and insure good coverage. Hold nozzle about one foot away while spraying. Apply until hair is thoroughly moist. Do not spray directly in eyes or mouth of animal. Do not repeat treatment for 24 hours. To reduce reinfestation, thoroughly spray the animal's bedding and other resting areas until the surface is slightly moist. Treat these areas at the rate of two seconds per linear foot at a distance of 12 to 18 inches. Also spray the entire inside surface of dog houses thoroughly spraying floors, crack, crevices, and bedding. Hit as many insects directly as possible. Do not repeat treatment for 24 hours. Do not use on puppies or kittens under 12 weeks. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals

To Protect Ornamental Plants in noncommercial greenhouses and lathouses, homes, commercial and institutional buildings, and for indoor landscaping: Spray directly on insects as directed below to kill aphids, exposed thrips, whiteflies, leafhoppers, plant bugs, spittlebugs, two-spotted mites, red spider mites, spider mites, cutworms and slugs on the following plants:

6/10

Directions for use continued:

African violet	coleus	asparagus fern	Jacob's ladder	pothos
ageratum	gold-dust croton	fuchsia	lilac	rose
amwranth	cup flower	geranium	lobellia	snapdragon
false aralea	daisy	heliotrope	marigold	silver vase
begonia	daphne	hibiscus	mayten tree	toyon
Christmas cactus	dracaena	impatiens	nasturtium	verbena
chrysanthemum	dumbcane	ivy	petunia	umbrella tree
cockscorb	fatshedera	jade	philodendron	wax plant
				zinnia

Contact Spray for pests listed above: Hold container 18 inches from the infested plants. Spray in short bursts from one to three seconds or spray at the rate of one linear foot of growth per second. Direct spray to all sides covering foliage until slightly moist. Avoid heavy wetting. Repeat as necessary to obtain control. If application is made indoors, remove infested plants from living and eating areas prior to spray applications. Spray several plants at once and in an area not to likely to come in contact with pets, children or food. When leaves dry, plants can be returned to original position. If plant removal is not possible, remove or cover exposed food and food handling surfaces, including utensils. Shut off ventilating systems, air ducts and air conditioning equipment. Avoid spray contact with surrounding surfaces in area. Leave the sprayed area after treatment and ventilate. Do not use treated plants for food or feed purposes.

Use with Caution on african violets, ash, azalea, reiger begonia, browallia, caenothus, camellia, columnea, cotoneaster, cyclamen, cypress, plumosa fern, gloxinia, impatiens, juniper, non-bearing mulberry and quince, oxalis, jade and primrose. As with all plants listed above, especially carnations, chrysanthemums and roses, there exists a wide variety of species and cultivars and, therefore, a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, spray a few plants as described above, wait 24 hours and check for possible injury.

Outdoors: For use around residential, commercial and institutional buildings, hospitals, gardens, noncommercial greenhouses, yards, patios, mobile homes, campers, boats, and swimming pools for controlling household and plant pests listed below:

To kill houseflies, mosquitoes, gnats (midges), deer flies, stable flies, flying moths (millers), fruit flies: Spray resting areas to reduce annoyance from these pests. Follow precautions listed above for ornamental plants. Hold container 18 to 24 inches from shrubs, bushes, and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 per cubic yard of space, while moving away from the treated areas. Treat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

Directions for use continued:

Wasps, hornets, yellow jackets, bees: Same as directed above for indoor use. APPLICATION SHOULD BE MADE LATE IN THE EVENING WHEN INSECTS ARE AT REST. Aim spray at nest openings in ground, bushes, and in cracks and crevices which may harbor nests, saturating nest openings and contacting as many insects as possible. Repeat as necessary.

Ants and fleas – Same as directed for indoor use.

Crickets, silverfish, earwigs, spiders, sowbugs, centipedes and millipedes - same as directed above for indoor use. TO PROTECT ORNAMENTAL PLANTS - Spray directly on insects as directed below to kill: **aphids, exposed thrips, whiteflies, leafhoppers, Japanese beetles, plant bugs, spittlebugs, two-spotted mites, red spider mites on:**

althea	forsythia	mimosa	pine	also:
white birch	hemlock	non-bearing citrus	pyracantha	mimosa webworm
cypress	holly	and cranberry	rhododendron	on
dogwood	locust	oak	spirea	mimosa
euonymous	maple	palm	spruce	
		viburnum	toyon	

To kill army worms, budworms, leaf beetles, gypsy moths, tussock moths, oakworms, sawflies, tent caterpillars, and weevils on the above plants; also **slugs** on roses; **spider mites** on chrysanthemums; **cutworms** on gladiolus and petunia. To kill **aphids, exposed thrips, adult whiteflies, leafhoppers, plant bugs and Japanese beetles on:**

African violets	chrysanthemum	fatshedera	ivy	philodendron
Ageratum	coleus	geranium	marigold	queen sago
Begonia	daisy	honeysuckle	nasturtium	roses
Broadleaf acacia	daphne	iris	peony	snapdragon
Tobira	zinnia			

Contact spray direction for above pests: Hold container 18 inches from the infested plants. Spray in short bursts from one to three seconds or spray at the rate of one linear foot of growth per second. Direct spray to all sides covering foliage until slightly moist. Avoid heavy wetting. Repeat as necessary to obtain control. For adult whiteflies and two spotted mites be sure to direct spray to contact these pests. Thoroughly cover the infested area. Use several repeat applications at two-day intervals if necessary.

Directions for use continued:

Use with caution on african violet, ash azalea, reiger begonia, browallia, caenothus, camellia, columnea, cotoneaster, cyclamen, cypress, plumosa fern, gloxinia, impatiens, juniper, non-bearing mulberry, flowering quince, oxalis, jade, primrose, and rose, including hybrid tea rose, there exists a wide variety of species and cultivars and , therefore, a wide range of susceptibility to injury. To ascertain possible phytotoxicity, spray a few plants as described above, wait 24 hours and check for possible injury.

Precautionary Statements**Hazards to Humans and Domestic Animals****CAUTION**

Cover and remove exposed foods. Avoid contamination of cooking utensils and food preparation surfaces. Remove pets and cover fish aquariums before spraying.

Use this Statement Only if Marketing for Dog & Cats**CAUTION**

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivities occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Certain medications can interact with pesticides. Consult a veterinarian before using on medicated animals.

FIRST AID**If In Eyes:**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

If on Skin or Clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting.

9/10

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to fish. For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in a cool, dry area inaccessible to children and pets. Keep container capped after use.

Disposal: Do Not Puncture or Incinerate!

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions.

NOTICE: This product contains a chemical known to the State of California to cause birth defects or other reproductive harm.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.

Important Note: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Sherwin-Williams Diversified Brands
601 Canal Road
Cleveland, OH 44113



5-1-03 10/10
NOTIFICATIONS

The Sherwin-Williams Company
601 Canal Road
Cleveland, Ohio 44113

April 16, 2003

Mr. Marion Johnson, PM 10
7505C
USEPA Headquarters
Ariel Rios Building
1200 Pennsylvania Avenue, N.W.
Washington, DC 20460

SUB: Notice of Label Change Relative to PR Notice 2001-6.
Disposal Instructions
EPA Reg. No.: 10900-63
876 Institutional Insecticide

Dear Mr. Johnson:

Enclosed please find an application for "Notification relative to PR Notice 2001-6, Disposal Instructions". I have enclosed a copy of the subject label for your review.

If you have any questions or need additional information please do not hesitate to contact me at (216) 515-8892.

Sincerely,

A handwritten signature in cursive script that reads 'Jamie Zajac'.

Jamie Zajac
Regulatory Affairs Specialist