

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH-567) WASHINGTON, D.C. 20460	EPA REGISTRATION NO.	DATE OF ISSUANCE
	10900-80	MAR 27 1985
NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	TERM OF ISSUANCE	
	NAME OF PESTICIDE PRODUCT	
Aqueous 30-3 Concentrate		

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Sprayon Products
 Division of Sherwin-Williams Company
 26300 Fargo Avenue
 Bedford Hts. OH 44146

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment:
 - a. Add the phrase "EPA Registration No. 10900-80."
3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

BEST AVAILABLE COPY

Timothy A. Gardner
 Product Manager (17)
 Insecticide-Rodenticide Branch
 Registration Division (TS-767)

Enclosure ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL  DATE 3-27-85

Lab #464

Sprayon's Aqueous 30-3 Concentrate

AN INSECTICIDE SPRAY CONCENTRATE

FOR USE IN FOOD PROCESSING AND STORAGE, MUSHROOM HOUSES, HOSPITALS, HOTELS, MOTELS, RESTAURANTS, OTHER INSTITUTIONS AND IN RESIDENCES

Authorized for use in all departments of official establishments operating under the meat, poultry, shell egg grading and egg products inspection programs.

PROVIDES RAPID KILL OF INSECTS WITHOUT LEAVING OBJECTIONABLE RESIDUES.

Food additive tolerances have been established for the insecticide components in this product. When this product is used as directed, allowable tolerance levels will not be exceeded.

Kills Gypsy Moth Caterpillars and Japanese Beetles

ACTIVE INGREDIENTS:

Pyrethrins.....	3.01
Piperonyl Butoxide, Technical.....	27.01
INERT INGREDIENTS:.....	100.01

*Equivalent to 24% (butylcarbityl)(6-propyl-piperonyl)ether and 6% related compounds.

PYRENONE - Registered Trademark of Fairfield American Corporation.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See reverse side for additional precautions.

EPA Reg. No. 10900- EPA Est. No. 10900-OH-1
10900- 10900-CA 1

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

NET CONTENTS

To prepare dilutions the concentrate should first be stirred or agitated well. Add required amount of concentrate to water. Blend thoroughly. This will produce a micro-emulsion.

For maximum effectiveness, a combination of localized surface application and space treatment is recommended.

AS A SURFACE SPRAY: For initial clean up or severe insect infestation, dilute at a rate of 1 part concentrate in 14 parts water (8.6 ounces per gallon). This dilution may be used in edible product areas of Federally Inspected meat and poultry plants. In other plants and for normal infestations dilute at a rate of 1 part concentrate in 29 parts water (4.3 ounces per gallon).

To control cockroaches, ants, silverfish, crickets, clover mites, cheese mites, ground beetles, rice weevils, confused flour beetle, rust red flour beetles, sawtoothed grain beetle, spider beetles, cigarette beetles, drug store beetles, meal worms, grain mites and cad Use a good sprayer adjusted to deliver a fine wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around corners of stored foods, around the base of machinery, behind shelves and drawers. Use in bookcases for silverfish. Spray ant nests and points of entry. If surface application only is to be used, spray floors, walls and other surfaces applying at a rate of 0.5 to 1.0 gallon per 750 square feet of surface.

To control carpet beetles, spray edges of walls and under carpeting and rugs. Make application to floor and baseboards. Spray directly into cracks, closets and infestations of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses especially particularly around tufts and seams. Take apart and spray in all joints. Treat bed moldings and floors. Repeat treatment as needed.

BATCH

To control fleas, and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. To kill fleas and ticks on cats and dogs, and to obtain protection against reinfestation, dilute at 1 part concentrate to 49 parts water (2.5 fluid ounces per 1 gallon) and wet the animal by dipping or spraying.

AS A SPACE SPRAY: Dilute at a rate of 1 part concentrate to 59 parts water (2 fluid ounces per gallon). Apply with mechanical or compressed air equipment (non-thermal) adjusted to deliver a fine mist. Close doors and windows and shut off ventilating systems. For rapid control of adult house flies, fruit flies, gnats, mosquitoes, shipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths: Direct spray at an upward angle distributing it uniformly through the entire area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for at least 10 minutes. Ventilate and sweep up dead insects before reoccupying.

For rapid kill of exposed or accessible stages of other insects named on this label; dilute 1 part concentrate in 5 - 79 parts water (4 to 25 ounces per gallon). Apply using conventional, mechanical or compressed air equipment (non-thermal), following directions for space spraying. Apply the above dilutions at a rate of from 1 to 10 ounces per 1000 cubic feet of space. When used in meat and poultry processing plants, a dilution of no less than 1 to 5 is acceptable and it must be dispersed from equipment adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns).

IN MUSHROOM PRODUCTION AND PROCESSING: To control adult mushroom flies, or fungus gnats, Phorids, Sciarids, and Cecidomyiids in growing houses. Dilute 1 part concentrate in 5 parts water (2.5 fluid ounces per gallon). Using mechanical or compressed air equipment (non-thermal) adjusted to deliver a fine mist, apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results refrigeration should be turned off. In packing and canning plants use at a rate of 1 ounce to 2000 cubic feet of space.

OUTDOORS: To kill gypsy moth caterpillars and adults, adult Japanese beetles and Eastern and forest tent caterpillars on contact: combine 1.0 fluid ounce of concentrate per 1 gallon of water. Apply to insure thorough coverage of upper and lower surfaces.

On Trees and Shrubs: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

On Roses and In Flower Gardens: Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES:

Store upright at room temperatures. Avoid exposure to extreme temperatures. In case of spill or leakage, suck up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

BEST AVAILABLE COPY

PRELIMINARY STATEMENTS HAZARD: TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Remove pets, birds and cover fish aquaria before spraying. Do not breathe spray mist. Wear mask or respirator of a type recommended by the U.S. Bureau of Mines to give adequate protection against this material. Wear protective clothing. Wash exposed skin surfaces with soap and water after handling and before eating or smoking. Wash all contaminated clothing with soap and water before reuse.

Do not apply as a space spray while food processing is underway. Except in federally inspected meat and poultry plants, when applied as a surface spray with care and in accordance with the directions and precautions given above, food processing operations may continue. Do not apply at any time as a undiluted space spray in edible product areas of federally inspected meat and poultry plants. Foods should be removed or covered before treatments. Thoroughly wash all food processing surfaces before reuse. All surfaces in processing areas must be washed and rinsed in potable water after spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams, ponds, tidal marshes, and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes.

CHEMICAL HAZARDS

Do not use or store near heat or open flame.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.