

FRONT PANEL

ARIZONA'S
BEST

DO IT YOURSELF
HOME PEST
CONTROL

CONTAINS DURSAM* INSECTICIDE FOR PROFESSIONAL
CONTROL OF INDOOR/OUTDOOR INSECT PESTS

KILLS CRICKETS, ROACHES, SPIDERS, SILVERFISH
ANTS, FLEAS, EARWIGS, BROWN DOG TICKS

ACTIVE INGREDIENTS:

Chlorpyrifos 0,0-diethyl 0-(3,4,5-trichloro-2-
pyridyl)phosphorothioate.....00.50%
Aromatic petroleum derivative solvent.....00.33%

INERT INGREDIENTS:.....99.17%

CAUTION: Keep out of reach of children
See directions and additional cautions
on side and back panels

NET CONTENTS 1 GAL. (3.785L)

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Recommended for use as a household insecticide for general pest control in and on buildings and structures and their immediate surroundings. Permitted areas of use include, but are not limited to: houses, garages, trailers, campers, storage sheds, apartment buildings, stores, warehouses, industrial buildings, schools, nursing homes, hospitals, restaurants and hotels.

INDOOR USE

This product is intended for application with a shaker or puffer container to hiding and runway areas and those places where pests are found. The amount to be applied will vary with the site, but should usually be in the range of 1/3 to 2/3 ounces of dust per square yard.

Cockroaches, Ants, Crickets, Silverfish and Firebrats: Thoroughly dust all areas where these pests crawl and hide, especially cracks and crevices around sinks, stoves, refrigerators and cabinets; behind baseboards; around doors and windows and in attic, basements and crawl spaces. For Ants, apply to ant trails and around doors and windows where they may enter premises.

Carpenter Ants: Dust in areas where carpenter ant nests are located. Nesting areas are usually found in woodwork, especially in areas where wood is located by water damage.

Brown Dog Ticks and Fleas in Buildings: Dust infested areas such as pet beds and resting quarters, nearby cracks and crevices, under the edges of rugs and floor coverings, between and under cushions of upholstered furniture and other places where these pests may hide. Old bedding of pets should be removed after treatment of pet area and replaced with clean fresh bedding. Additionally, ensure that pets are treated with an EPA approved product registered for pet use. **DO NOT TREAT ANIMALS OR THEIR BEDDING WITH THIS PRODUCT.**

Spiders: Dust along and behind baseboards, window and door frames, corners, pipes, storage localities, attics, basements, crawl spaces and other areas where spiders may hide.

DIRECTIONS FOR USE:

Wasps: It is best to treat wasp nests at night when insects are less active and have returned to the nest. Do not wear loose fitting clothing. Take necessary precautions to avoid stings. For best results, carefully dust near nest and check one or two days after treatment to ensure complete kill, and retreat if necessary. If possible, remove and destroy nest to prevent hatching of new insects. For ground nests, carefully dust entrance and surrounding areas. Retreat in one or two days if wasp activity is still present. Do not use this product to treat nests overhead.

Sowbugs, Pillbugs, Ground Beetles, Earwigs, Millipedes, Centipedes and Scorpions: Apply around doors, windows and other places where these pests may enter. Treat baseboards, storage areas, and cracks and crevices, especially in dark, musty areas where these pests may hide.

OUTDOOR USE

Perimeter Treatment: Use as a residual treatment for control of Ants, Brown Dog Ticks, Centipedes, Cockroaches, Crickets, Earwigs, Firebrats, Fleas, Ground Beetles, Millipedes, Pillbugs, Scorpions, Silverfish, Sowbugs, Spiders and Wasps. Make a residual treatment around windows and doors, porches, screens, eaves, patios, garages, under stairways and in crawl spaces and other areas where pests hide.

To help prevent invasion of buildings and structures by the above pests, treat a band of soil and vegetation 6 to 10 feet wide around and adjacent to the building or structure. Also treat the building foundation and crawl spaces where pests are active and may find entrance. Apply thoroughly and uniformly to the band area. Pay particular attention to ant hills and wasp nests in the band area. Wasp nests in trees and shrubs may be treated as noted above.

Imported Fire Ants: Apply 1/2 oz. of product to 2 gallons of water. Stir 1/2 oz. of product into 2 gallons of water. Pour a small amount of material around the perimeter of the mound. Apply the remaining volume directly on the mound so as to penetrate insect galleries, using 2 gallons of mixture on mounds 8 inches or larger in diameter. Less mixture can be used on smaller mounds; however, thorough penetration of the insect galleries must be achieved. A second treatment may be necessary if complete control is not obtained.

Do not disturb the mound prior to application. If possible make applications after a recent rainfall. Applications may not be as effective during hot, dry periods.

To Make A Dust Application: Mix 1/2 oz. of product with 1/2 cup of dry material between 1/2" and 1" in size. Apply the mixture to the perimeter of the area surrounding the mound. Apply the mixture to the center area where the mound is active. Repeat applications as needed.

INDOOR USE (Cont.)

FLEAS & TICKS IN CARPETS

easy application. Apply powder to carpet surfaces. Wait 30 minutes. Vacuum powder off carpet surfaces only. Use vacuum cleaner and sweep with broom. Dispose of vacuum in trash. Keep pet off carpet until 24 hours after carpet has been treated. Do not vacuum carpet before treatment. This product will kill Adult Fleas, Flea Larvae, Flea Pupae, Silverfish & Spiders.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION:

Harmful if swallowed inhaled or absorbed through the skin. Do not breathe dust. Do not allow to contact skin, eyes or clothing.

Wash hands and exposed skin before eating, drinking or smoking and after handling. Wash all contaminated clothing thoroughly before reuse.

Do not apply to humans, animals, clothing or bedding.

Do not use in food preparation areas, nurseries, day care centers, or other areas where children play.

Do not contaminate feed or feed products or feed preparation surfaces, dishes, kitchen utensils and food containers.

ENVIRONMENTAL HAZARD

This product is toxic to fish and aquatic life. Do not discharge directly to water. Do not discharge to sewer. Dispose of waste properly.

This product is toxic to birds, bees, butterflies, and other beneficial insects. Do not apply to crops. Do not apply to ornamental plants. Agriculture Use Only.

STORAGE AND DISPOSAL *12 PT*

Do not store in children's reach. Store product in original container. Do not use if container is damaged or if contents are spilled. Do not use if product is expired.

For more information, contact the manufacturer.

It is a violation of the law to sell or give away any insecticide without a label. The label must be printed in plain English and must contain the following information:

AS A BAIT FOR THE CONTROL OF THE HOUSEHOLD CRICKET

NOTE
Do not use this product where food crops are raised, such as corn, wheat, radishes, etc. Do not use where children are likely to play or their parents are likely to be present.

Apply this product to the soil at a rate of 1 lb. per 100 sq. ft. (1 qt. / 100 sq. ft.). Sprinkle over the soil in the area where these insects are seen or are likely to be seen after it is consumed or as needed for continued control. Apply approximately 30 lbs. per acre.

Apply in late afternoon when the soil is moist where insects have been seen or are known to exist.

MOLE CRICKETS in lawns, golf courses and Residential Turf - For control of Mole Crickets apply this granular product at the rate of 3 1/2 pounds per acre or approximately 1 1/2 lbs. per 1000 sq. ft. over the area using a mechanical spreader. Apply in late afternoon and moist soil in areas known to be infested with these insects. Repeat application is about one week for best control.

THE BEST TIME TO CONTROL THE SOUTHERN MOLE CRICKET

Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
	2 /	1 /									

NOTE - #2 indicates the best time to control the Adult Mole Crickets.
#1 indicates the laying time during the summer