

BIONICS

Bionite

Disinfectant & Sanitizer



4-4-72
6962-34

10910-5

Directions for Dilution

DILUTE WITH WATER AS DIRECTED
DO NOT
MIX WITH SOAP OR ANIONIC DETERGENTS

ARTICLES TO BE SANITIZED SHOULD
FIRST BE WASHED AND RINSED

For 200 ppm Solution
Add and Stir 2.5 oz. BIONITE
in 10 gallons of water.

For 400 ppm Solution
Add and Stir 5 oz. BIONITE
in 10 gallons of water.

For 800 ppm Solution
Add and Stir 10 oz. BIONITE
in 10 gallons of water.

This product fulfills the criteria of Appendix F
of the Grade "A" Pasteurized Milk Ordinances
1965 Recommendation of the U.S. Public
Health Service in waters up to 550 ppm of
hardness calculated as CaCO_3 when tested by
the AOAC Germicidal and Detergent Sanitizers
— Official Method.

E.P.A. Reg. No. 10910-5

Net contents U.S. gallons

BIONICS INDUSTRIES 11250 W. ADDISON ST. / FRANKLIN PARK, ILLINOIS 60131 / (312) 455-8400

Los Angeles - 90248
(213) 327-6600

Dallas - 75220
(214) 350-4046

New York - 11378
(212) 894-5151

INSTRUCTIONS

EATING & DRINKING ESTABLISHMENTS

DIRECTIONS FOR USE:

1. Scrape and prewash utensils and glasses whenever possible.
2. Wash with a good detergent or a compatible cleaner.
3. Rinse with clean water.
4. Sanitize in a solution of 2½ oz. of BIONITE to 10 gallons of water (200 ppm). Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drain-board to air dry.

ACTIVE INGREDIENTS:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈)	
dimethyl benzyl ammonium chlorides	5.0%
n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl	
ethyl benzyl ammonium chlorides	5.0%
INERT INGREDIENTS:	90.0%
	100.0%

WARNING: KEEP OUT OF REACH OF CHILDREN.

Causes eye and skin irritation. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food. FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, do not induce vomiting. Drink large quantities of fluid and call physician. Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

APPROVED UNDER THE REGULATIONS OF THE WISCONSIN STATE BOARD OF HEALTH

Milk Farms, Creameries and Food Processing Plants: First clean utensils and equipment with a good detergent, rinse well and then sanitize with solution 200 ppm by immersion for 2 minutes. Use same dilution for flank and udder wash prior to milking. Before re-use of equipment coming in direct contact with food, a rinse with clear running water is recommended. Clean floors, walls and storage areas with a good detergent, rinse well, then wipe or mop with solution 400 ppm or spray with solution 800 ppm.

Sanitizing Shell Eggs: BIONITE can be used to sanitize shell eggs. When BIONITE-sanitized eggs are to be cartoned no rinse is required. When BIONITE sanitized eggs are to be immediately broken for use in the manufacture of egg products, then the eggs must be subjected to a thorough potable water rinse.

MADE & PRINTED IN U.S.A.

© 1971 CHEMTRUST INDUSTRIES CORP.

B-14-2