



**NET CONTENTS  
10 POUNDS**

**INGREDIENTS**

ACTIVE INGREDIENTS	
MANEB Manganese ethylene bisdithiocarbamate	50.00%
HEXACHLOROBENZENE	10.00%
INERT INGREDIENTS 40.00%	

EPA REG. NO. 10912 ■ 1

**Directions for Use**

- WHEAT Use rate of 1 oz. per bushel
- BARLEY OATS RYE Use rate of 2 oz. per bushel
- FLAX Use rate of 4 oz. per bushel
- CORN and SOYBEANS Use rate of 2 1/4 oz. per bushel

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN.**  
**Harmful if swallowed. Avoid inhaling fumes. This material may cause skin irritation. Wash hands and face thoroughly after using. Do not store near heat or open flame. Use with adequate ventilation. Keep container closed. Do not use treated seed for food, feed or oil purposes. Avoid contact with skin, eyes or clothes. See other precautions on the right side panel.**

**SEED DISINFECTANT AND INSECTICIDE  
FOR DRILL OR PLANTER BOX APPLICATION**

**CAUTION**

**NOTE:** Do not use treated seed for food or feed purposes.  
 This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of water. Apply this product only as specified on this label.  
 Keep out of reach of children when not in use. If Manganese ethylene bisdithiocarbamate and Hexachlorobenzene becomes wet under agricultural conditions, its effectiveness may be reduced and it may also produce a harmful vapor. Store in a cool dry place.

**DESTROY EMPTY CONTAINERS.** Destroy empty containers by puncturing, crushing, and burning in rear crop land.

**NOTE:** Treat seed in rear crop land as follows:

- 1. DISINFECTED SEED
- 2. CONTAMINOUS TO ANIMAL AND MAN
- 3. DO NOT USE FOR FOOD, FEED OR OIL

- Mix THIHEX thoroughly into the seed.
1. Placing and leveling one-half of the seed in the drill or planter box and sprinkle one-half the total THIHEX required for the full planter, uniformly on or under the surface of the seed. Mix thoroughly with a stick or paddle, then fill the box with seed and sprinkle the remaining one-half of the powder on or under the surface and mix again until all streaking has disappeared and coloring of seed appears uniform. **DO NOT MIX WITH HANDS.**
  2. Dribble the required amount of THIHEX into each bushel of seed as it is poured into the drill box and mix with a paddle until uniform.
  3. For Slurry Machines: Adjust to apply one half pint of slurry per bushel of seed and use 2 lbs. of seed treatment per gallon of water. If machine applies more or less slurry, adjust the amount of seed treatment to apply 2 ounces per bushel of seed. For Flax seed apply 4 ounces per bushel.
- Avoid very deep seeding and exceptionally early sowing under poor conditions for maximum benefits. Use only recommended rates. Seed should be well cured, dry and clean before treatment.

**Product Information**

THIHEX is a special seed treatment formulation of Manganese ethylene bisdithiocarbamate (Maneb) and Hexachlorobenzene (HCB) which, when used in accordance with Directions for Use, protects crops against certain soil borne diseases such as ~~stink bug~~ ~~and other~~ ~~soil~~ ~~borne~~ ~~diseases~~ such as smut, covered and false looge, root of barley, and bunt of eye. It also gives temporary protection to cereal crops, corn and soybeans, against certain soil borne diseases such as root rot and seedling blight.

*Aug 19, 1974*  
 10102-3