

PRECAUTIONARY STATEMENTS

Hazards to humans and domestic animals.

CAUTION

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling and before smoking or eating. Avoid contamination of food and feed.

STATEMENT OF PRACTICAL TREATMENT

If on skin, remove by washing thoroughly with mild soap and water. If in eyes, wash with copious amounts of water and get medical attention. If swallowed, drink promptly a large quantity of milk, egg whites, or gelatin solution. See physician immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Application rates: Use lower rates with normal infestation and higher rates with severe infestation.

STORAGE AND DISPOSAL ^{Upoints}

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinseate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL:

Household: Do not reuse empty container.

Wrap and put in trash collection.

Plastic: Triple rinse (or equivalent) and offer for recycling or recondition, or dispose of in a sanitary landfill or by incineration if permitted by State and local authorities.

ACCEPTED
with COMMENTS
EPA Letter Dated:

FEB 22 1983

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
a pesticide registered under EPA Reg. No.

FS-6

FLOWABLE SULFUR

| | |
|------------------------|---------------|
| Active Ingredient | |
| Sulfur..... | 52.0% |
| Inert Ingredients..... | 48.0% |
| | <u>100.0%</u> |

This product contains 5.0 pounds Sulfur per gallon.

KEEP OUT OF REACH OF CHILDREN - 12 points

CAUTION - 12 points

See side panel for additional Precautionary Statements.

DIRECTIONS FOR MIXING: BEFORE USING, STIR UNTIL SMOOTH

Pour recommended amount in partially filled spray tank. Keep agitator running during filling and spraying operation. DO NOT ALLOW MIXTURE TO STAND. Failure to maintain agitation will cause FS-6 Flowable Sulfur to settle and may necessitate manual stirring to redisperse.

The strong adhesive properties of FS-6 Flowable Sulfur act as a sticker on the plant, and the sticking characteristic necessitates the flushing of equipment with water after each day's use. Sulfur in any form is corrosive material. TO REDUCE THE EFFECT, EQUIPMENT SHOULD BE FLUSHED DAILY.

APPLICATION INSTRUCTIONS

Fruit tree dosage rates are given as pints of FS-6 Flowable Sulfur per 100 gallons of water. Wet all foliage. When there is a question as to rate of application, use the higher rate.

Apple, Pear - Scab

- Pre-bloom through Calyx sprays 1 1/2 to 3 1/2 pints
- Cover sprays 1 to 2 pints
- Powdery Mildew:
- Pre-bloom through Calyx sprays 1 1/2 to 3 1/2 pints
- Cover sprays 2 pints

Cherry, Plum, Prune: Brown rot, Leafspot:

- Pink and bloom sprays 2 1/2 to 3 pints
- Petal fall shuck and cover sprays 2/3 to 1 3/4 pints

Citrus: Rust Mite, Clover Mite:

1 to 4 pints

Apply sprays November thru May

Use 100 gallons of water per acre for thorough coverage. Apply at first sign of infestation and repeat at 2 week intervals. For control of rust mite, apply at first sign of infestation and repeat at 2 week intervals.

Use 100 gallons of water per acre for thorough coverage. Apply at first sign of infestation and repeat at 2 week intervals. For control of rust mite, apply at first sign of infestation and repeat at 2 week intervals.

Apple, Pear - Powdery Mildew, Brown Spot, Leaf Spot:

- Pink and bloom sprays 1 to 3 pints
- Petal fall, shuck and cover sprays 2/3 to 1 3/4 pints

Citrus - Powdery Mildew:

Use 3 to 4 pints in 100 gallons water per acre by air. Apply at first sign of infestation and repeat at 2 week intervals.

Strawberries - Powdery Mildew:

Use 1 pint per acre in sufficient water for thorough coverage. Apply at first sign of infestation and repeat at 2 week intervals. For strawberries that will be canned, harvest a portion of sulfur well before harvest to reduce risk with local sulfur dioxide conditions.

Gooseberries - Powdery Mildew:

Use 1 to 2 pints per acre in sufficient water for thorough coverage, depending on size of bush, or so to wet all or all foliage.

Plum - Powdery Mildew:

Apply 1 to 2 pints per acre in sufficient water for thorough coverage, or apply 1/2 to 1 pint per bush in sufficient water for thorough coverage.

Plum - Rust:

Apply 1 to 4 pints per acre in sufficient water for thorough coverage. For control of rust, apply at first sign of infestation and repeat at 2 to 3 week intervals. Do not apply more than 4 applications.

Plum - Leaf Spot:

Use 100 gallons of water per acre for thorough coverage. Apply at first sign of infestation and repeat at 2 week intervals.

Application Instructions Continued

2-4 pints per acre in the Pacific Northwest and 2-8 pints per acre in Arizona, Colorado, Nebraska, Wyoming, New Mexico, Texas, and Utah. Repeat applications should be made at 10 to 30 day intervals or as necessary throughout the season. Under most disease conditions only one or two applications will be needed.

PRODUCT INFORMATION

NOTE: Some fruits and other plants are susceptible to injury from sulfur under climatic conditions. The user is advised not to use sulfur on any crop unless local use has proven that sulfur is safe in the locality.

During periods of high temperature sulfur may burn foliage and fruit. Do not make sulfur applications at excessively high temperatures, ie 90° F. or above. Do not use FS-6 Flowable Sulfur with oil or within four weeks of an oil application.

When FS-6 Flowable Sulfur is used with arsenicals, lime should be added to prevent plant injury.

Consult your State Agricultural Experiment Station or State Extension Service Specialist for complete details on the spraying program best suited to your local conditions.

THIS PRODUCT FREEZES at 32° F but usually upon thawing is still useable.

NOTICE

Use of this material in a manner other than in accordance with this label may cause serious plant or crop injury, excessive residues and unsatisfactory control.

Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use; but neither this warranty or any other warranty of merchantability of fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions not reasonably foreseeable to the seller, the buyer assumes the risk of any such use.