

disp

ACCEPTED
with COMMENTS
in EPA Letter Dated:

NOV 29 1982

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

10912-28

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

May be fatal if swallowed. May be absorbed through skin. Harmful if inhaled. Causes eye and skin irritation. Do not get in eyes, on skin or on clothing.

Wash hands, arms, and face thoroughly with soap after handling and before eating or smoking. Routinely shower or bathe after working and wash all clothing with soap and hot water before re-use. During exposure in seed treatment and loading operations wear goggles, clean synthetic rubber gloves, protective clothing, and a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR part 11.

Avoid contamination of feed and foodstuffs. Do not store with food. Do not store with feed or forage to be fed in dairy animals or to animals being finished for slaughter.

Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time. Seed commercially treated with this product must be labeled in accord with all applicable requirements of the Federal Seed Act.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Exposed treated seeds may be hazardous to birds and other wildlife. Dispose of all excess treated seeds by burial away from bodies of water. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTION FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Wheat — Use rate of 3.0 oz. per bushel.

Barley, Oats, Rye — Use rate of 4 oz. per bushel.

Flax — Use rate of 8 oz. per bushel.

Corn and Soybeans — Use rate of 5½ oz. per bushel.

Mix Trinox Liquid thoroughly into the seed.

(1) Placing and leveling one-half of the seed in the drill or planter box and apply one-half of the total Trinox Liquid required (for the full planter box) uniformly on or under the surface of the seed. Mix the remaining one-half of the Trinox Liquid on or under the surface and mix again until all streaking has disappeared and coloring of seed appears uniform. DO NOT MIX WITH HANDS.

(2) Dribble the required amount of Trinox Liquid into each bushel of seed as it is poured into the drill box and mix with a paddle until uniform.

Avoid very deep seeding and exceptionally early sowing under poor conditions for maximum benefits. Use only recommended rates. Seed should be well cured, dry and clean before treatment.

LIQUID TRINOX LX SEED TREATMENT

ACTIVE INGREDIENTS:

Maneb (Manganese ethylene bisdithiocarbamate)	25.00 %
Lindane (Gamma Isomer of Benzene Hexachloride)	9.375 %
INERT INGREDIENTS	65.625 %
	100.00 %

KEEP OUT OF REACH OF CHILDREN

WARNING

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person or convulsing person.

IN CASE OF EYE CONTACT: Immediately flush eyes with running water for at least 15 minutes. Call a physician.

IN CASE OF SKIN CONTACT: Wash the affected area with soap and warm water.

IF INHALED: Remove victim to fresh air. Apply respiration if indicated.

Net Contents 2½ Gallons

EPA Est. No. 10912-MN-1

EPA Reg. No. 10912-28

Manufactured by Haynes Chemical Company
P. O. Box 30 — East Grand Forks, MN 56721

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool place. Do not store in direct sunlight. Do not allow to freeze.

PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by state and local authorities.

PRODUCT INFORMATION

Trinox Liquid is a special seed treatment formulation of Manganese ethylene bisdithiocarbamate (Maneb) and Gamma Isomer of Benzene Hexachloride (Lindane) which, when used in accordance with Directions for Use, protects crops against certain soil-borne diseases such as stinking smut of wheat, oat smut, covered and false loose smut of barley, and bunt of rye. It also gives temporary protection to cereal crops, corn and soybeans against certain soil-borne diseases such as root rot, seedling blight and for temporary protection of seeds and seedlings against wireworms.

Trinox Liquid is especially designed for on-the-farm application as a Drill Box Treatment.

WARRANTY STATEMENT

Haynes Chemical Company warrants that the product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label. Haynes Chemical Company neither makes, nor authorizes, any agent or representative to make any other warranty of fitness or merchantability, guarantee or representation, expressed or implied, concerning this material.

