

## DIRECTION FOR USE

Dormant and delayed dormant application. Almonds, apricots, apples, cherries, peaches, pears, prunes. Apply 5-8 gallons oil per acre for control of European Red Mite, Brown Almond Mite, two-spotted Mite, Pacific Mite, Pear Psylla, McDaniel Mite, Strawberry Mite, Plum Nursery Mite, Wilamette Mite, Pear Rust Mite, Peach Silver Mite, San Jose Scale, and Lecanium Scale.

**DILUTION RATE:** Apply 2 gallons of Superior "98" Spray Oil per 100 gallons of water.

Fill spray tank with 1/2 the required amount of water. Start agitator. Add total amount of spray oil. Continue operation of agitator while adding balance of water and while spraying.

Spray oil may be used as a carrier for appropriate Insecticides and Fungicides which are Registered for use on deciduous fruits and nuts and which can be applied to give a uniform spray. Consult your local Agricultural Extension Service.

**GENERAL DIRECTIONS:** Superior "98" is a straight-cut narrow boiling range petroleum oil designed specifically for use on deciduous fruit trees.

Apply so as to give total coverage to all parts of tree. Thorough drenching of trunks limbs, twigs and leaves is essential for maximum insect control.

USDA Reg. No. 10914-1

NET CONTENTS \_\_\_\_\_ GALS.

# FEED SERVICE SUPERIOR "98" SPRAY

## NEW SUPERIOR SPRAY OIL FOR DECIDUOUS FRUITS

Active Ingredient	
Mineral Oil	98.0%
Inert Ingredient	2.0%
	<hr/>
	100.0%

Unsulfonated Residue  
(ASTM D-483) 93%

Petroleum Oil Classification: Unclassified

Mfgd. with PGSO-2 Base Oil

**STRAIGHT-CUT NARROW RANGE OIL**

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

Harmful if Swallowed. Do Not Induce Vomiting.

Call a Physician Immediately.

Avoid Inhalation of Vapors  
and Spray Mist

Manufactured By:

# FEED SERVICE INC.

Caldwell, Idaho

## CAUTION

**READ ENTIRE LABEL:** Use strictly in accordance with label cautions, warnings, and directions; and in conformity with Federal and State regulations.

**DO NOT APPLY** spray oil without consulting your State Agricultural Extension Service.

**DO NOT COMBINE WITH SULFUR SPRAYS** and **DO NOT APPLY** spray oil within 30 days before or after an application of sulfur.

**DO NOT MIX** with insecticides or fungicides incompatible with oil. Consult your State Agricultural Extension Service for specific compatibilities.

**DO NOT APPLY** to trees weakened by disease, drought, drying winds, or high nitrogen applications.

**DO NOT USE SUPER SPRAY** when temperatures are above 90° Fahrenheit.

## KEEP OUT OF REACH OF CHILDREN

Harmful if Swallowed—Do Not induce Vomiting

**CONDITIONS OF SALE:** 1. FEED SERVICE, INC. (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty of **FITNESS** or of **MERCHANTABILITY**, guarantee or representation, express or implied, concerning this material. 2. Critical and unforeseeable factors beyond the manufacturer's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above) resulting from handling, storage, and use of this material.