

WEED-OUT, a fertilizer & weed control for blade grass lawns only

ACTIVE INGREDIENTS	
*Dimethylamine salt of 2-2 Methyl 4 Chlorophencky Propionic Acid	.576%
**Omethylamine salt of 2.4 Dichlorophenoxyacetic acid	878%
the Compathy against gamest by graphy at the fact there is a consider a set	744
NERT INGREDIENTS	98 492%
, * ita	100 000%
*Equivalent to 2-2 Methyl 4 Chiorophenoxy, Propionic Acid	476%
"Equivalent to 2.4 Dichlorophenoxyacetic acid	'29 %
***Edurvalent to 3.6 Dichloro o anisic acid	()45%

24 lbs. covers 5000 sq.ft.

E P A REG NO 10942 2

•

CAUTION: KEEP OUT OF REACH OF CHILDREN AND PETS

AVOID CONTACT WITH SKIN, EYES OR CLOTHING. WASH ANY EXPOSED SKIN THOROUGHLY.

WASH ANY EXPOSED SKIN THO HARMFUL IF SWALLOWED.

KEEP CHILDREN AND PETS OFF TREATED AREA UNTIL GRASS IS ALLOWED TO DRY. AVOID CONTAMINATION OF FEED AND FOODSTUFFS

VAPORS FROM THIS PRODUCT MAY INJURE SUSCEPTIBLE PLANTS I'I THE IMMEDIATE VICINITY.

AVOID DRIFT TO DESIRABLE PLANTS.

DO NOT CONTAMINATE WATER USED FOR IRRIGATION OR DOMESTIC PURPOSES.

AVOID APPLICATION ON SHRUBS, FLOWERS, FRUIT, VEGETABLES, DICHONDRA, OR ANY DESIRABLE BROADLEAF PLANTS OR AREAS UNDERLAID BY ROOTS OF DESIRABLE TREES AND SHRUBS.

MAY CAUSE INJURY TO ST. AUGUSTINE GRASS, BENTGRASS AND CARPETGRASS.

DO NOT APPLY TO NEWLY SEEDED AREA.

DO NOT RESEED UNTIL AT LEAST 120 DAYS AF' ER APPLICATION.

CLEAN AND WASH APPLICATION EQUIPMENT THOROUGHLY AFTER USE.

DO NOT STORE NEAR FERTILIZER, SEEDS, INSECTICIDES, OR FUNGICIDES.

DO NOT REUSE CONTAINER. DESTROY BY BURYING IN A SAFE PLACE AWAY FROM WATER SUPPLIES. DO NOT BURN.

THIS PRODUCT IS A COMBINATION OF PROVEN EFFECTIVE SYSTEMIC ACTION HERBICIDES IN CONJUNCTION WITH A MILD FERTILIZER. THE FOLLOWING IS A PARTIAL FIST OF WEEDS CONTROLLED BY ROPER USE OF THE PRODUCT.

· ·			
F REDSTRAM	FLET TO TK	M STARD	Conference of the fi
C NOMEED	Att 1.150 A. 1	14, 5	* MAG* A; F5
THE ACK MEDIC	ME AL A	MARKET STATE A	S. M. AER
ARPET MEED	++E 7. ESI T	F 34115	JPC TTED SPURGE
THICK WEED IN SAME A	MAOTAFED	P. AT. TAIN	THETLE
" "HI" HAFED Nº USE EAR	LANGSGLARTER	F RS. Att	All Districts
CEA	MILECAD LAND A	MA IMPED	Vi. L. JAPI
DANDEL-CA.	MCRUNGS	SHEFHERD'S P HSE	WILD LETTUCE
DICHUIT COS			

INSTRUCTIONS FOR USE: (FOR BLADE GRASS LAWNS ONLY)

READ AND FOLLOW ALL INSTRUCTIONS CAREFULLY

WHEN TO APPLY:

APPLY ANYTIME THROUGHOUT THE YEAR WHEN ANY BROADLEAF WEEDS ARE PRESENT: MOST EFFECTIVE WHEN WEEDS ARE GROWING ACTIVELY. A REPEAT APPLICATION MAY BE MADE IN FOUR WEEKS IF NECESSARY.

HOW TO APPLY

NOTE: DO NOT APPLY ON A WINDY DAY, OR WHEN AIR TEMPERATURE EXCEEDS 90 DEGREES

- 1. MEASURE AREA TO BE TREATED (FOR EXAMPLE 50 BY 100 FEET EQUAL 5000 SQ FT.) SPREAD CONTENTS OF THIS PACKAGE EVENLY OVER 5000 SQ FT. OF LAWN AREA LAWN SHOULD BE CUT 2 OR 3 DAYS BEFORE APPLICATION AND WATERED THOROUGHLY THE DAY BEFORE APPLICATION.
- 2 WATER LAWN LIGHTLY JUST BEFORE APPLICATION, SO MATERIAL WILL ADHERE TO LEAVES DO NOT WATER AGAIN FOR AT LEAST 24 HOURS.
- 3. DO NOT OVERLAP

NOTE: EVEN A SLIGHT OVER-APPLICATION MAY CAUSE TEMPORARY DISCOLORING OF SOME VARIETIES OF GRASS WITH NORMAL CARE. RECOVERY SHOULD OCCUR IN 2 TO 3 WEEKS

APPROXIMATE SPREADER SETTINGS USE DROP TYPE SPREADER ONLY.

IF YOU WALK SLOWER THAN NORMAL USE A HALF NUMBER SMALLER, IF YOU WALK FASTER THAN NORMAL USE A HALF NUMBER LARGER.

BANDINI

4¹2

SCOTT (MODEL 35)

5¹4


PLEASE CHECK WITH DEALER FOR PROPER SETTING OF OTHER SPREADERS.

CAUTION: ALWAYS SHUT OFF SPREADER WHEN FILLING, MAKING SHARP TURNS OR STOPPING. AVOID SPILLING.

PRODUCT OF BANDINI FERTILIZER COMPANY LOS ANGELES, CALIFORNIA 90023, U.S.A.

Jan 23, 1973

10942-2





WEED-OUT. a fertilizer & weed control for blade grass lawns only

****Dimethylemine salt of Dicambe (3, 6-Dichloro-o-anielo acid)

*Equivalent to 2-(2-Methyl-4-Chiorophenoxy) Propionic Acid
**Equivalent to 2, 4-Dichlorophenoxyscatic acid

24 lbs. covers 5000 sq.ft.

E. P. A. REG. NO. 10942-2

CAUTION KEEP OUT OF REACH OF CHILDREN AND PERS

INSTRUCTIONS FOR USE I FOR BLADE GRASS LAWNS OF THE